




Volume 40 • Number 12 • December 1978

Marine Fisheries REVIEW

National Oceanic and Atmospheric Administration • National Marine Fisheries Service



Seafood . . .

. . . nutrition.

Vitamins

Marine Fisheries Review

Vol. 40, No. 12
December 1978

CONTENTS

Articles

- 1 Composition of the Edible Portion of Raw (Fresh or Frozen) Crustaceans, Finfish, and Mollusks. IV. Vitamins
Virginia D. Sidwell, Audrey L. Loomis, Pauline R. Foncannon, and David H. Buzzell
- 17 Review of Oxygen Depletion and Associated Mass Mortalities of Shellfish in the Middle Atlantic Bight in 1976
Frank W. Steimle and Carl J. Sindermann

Departments

- 27 NOAA/NMFS Developments
- 30 Foreign Fishery Developments
- 32 Publications
- 34 Index, Volume 40

Cover: The vitamins in fishes (see article on page 1) are reflected in the illustration by Harold Spiess.

U.S. DEPARTMENT OF COMMERCE
Juanita M. Kreps, Secretary

NATIONAL OCEANIC AND
ATMOSPHERIC ADMINISTRATION
Richard A. Frank, Administrator
Terry L. Leitzell, Assistant
Administrator for Fisheries

National Marine Fisheries Service



Marine Fisheries Review is published monthly by NMFS Scientific Publications Staff, Room 450, 1107 N.E. 45th St., Seattle, WA 98105.

Publication of material from sources outside the Service is not an endorsement. The Service is not responsible for the accuracy of facts, views, or opinions of these sources.

Although the contents have not been copyrighted and may be reprinted freely, reference to source is appreciated.

The Secretary of Commerce has determined that the publication of this periodical is necessary in the transaction of the public business required by law of this Department. Use of funds for printing this periodical has been approved by the Director of the Office of Management and Budget through May 31, 1980.

Managing Editor: W. Hobart

For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, DC 20402. Price \$1.10 (single copy). Subscription price: \$12.75 a year, \$15.95 a year for foreign mailing. Controlled circulation postage paid at Tacoma, Wash.

Composition of the Edible Portion of Raw (Fresh or Frozen) Crustaceans, Finfish, and Mollusks. IV. Vitamins

VIRGINIA D. SIDWELL, AUDREY L. LOOMIS,
PAULINE R. FONCANNON, and DAVID H. BUZZELL

ABSTRACT—This report summarizes data from 157 publications referring to the vitamin content of commonly eaten seafoods. Included are references to vitamins A, D, and E, choline, ascorbic acid, inositol, biotin, thiamin, riboflavin, niacin, pyridoxine, folic acid, vitamin B₁₂, and pantothenic acid.

INTRODUCTION

In Part I, Sidwell et al. (1974) described the data bank being established at the Southeast Fisheries Center College Park Laboratory¹ of the National Marine Fisheries Service, NOAA, College Park, Md. The objectives of the data bank project are: 1) To develop a comprehensive, systematic data retrieval system containing available information on the chemical and nutritional composition of fish and fishery products; 2) to publish information on the nutrients found in fishery products based on a wide survey of the literature and 3) to point out aspects of chemical composition of fish which need further investigation. Part I also summarized the data on the content of protein, fat, moisture, ash, carbohydrate, energy (calories), and cholesterol in commonly eaten seafoods.

In Part II, Sidwell et al. (1977) reviewed the data on the concentration of sodium, potassium, calcium, phosphorus, chlorine, and magnesium found in edible portions of seafood.

In Part III, Sidwell et al. (1978) as-

sembled the data from literature on the concentrations of various microelements (trace elements) in seafood.

This paper, Part IV, summarizes the data on the various vitamins in fish and shellfish muscle. There is a need in the medical community for such information, e.g., in the evaluation of nutritional aspects and in the calculation of special diets. Industry can use the data to encourage the use of nutritional labeling. And the nutrition-conscious consumer wants to know the nutritional aspects of fish.

RESULTS AND DISCUSSION

In this report we assembled, from 157 references, data on the vitamins present in the flesh of 140 different species of aquatic animals.

The averages and ranges for each vitamin were calculated from the data reported by various investigators. These data were in the form of an average based on multiple determinations, or were reported as individual values from which an average could be calculated. Regardless, only averages were used to calculate the overall average, standard error of the mean, and range appearing in Tables 1 and 2. With the aforementioned statistics is the number of aver-

ages used to make the statistical calculations.

A limited number of analyses are available for many species of finfish or shellfish listed in Tables 1 and 2. A number of commonly caught species have not been analyzed for vitamin content. Wherever the information for a specific fish and/or vitamin is limited, the recorded value should be regarded only as an estimate of what can be expected. More data are necessary to obtain a value that has a higher degree of reliability.

As noted in Tables 1 and 2, the range of values for each of the vitamins in the flesh of the same species of animal is quite large. A portion of this variation within each species is undoubtedly associated with seasonal and biological differences, e.g., size of the animal, its age, sex, degree of sexual maturation, and method of handling the fish after it left the water and before the flesh was analyzed for the various vitamins. Also, some of the variation may be related to the technique used in the preparation of the sample before it is analyzed or the method of analysis used by the investigator.

Virginia D. Sidwell, Audrey L. Loomis, Pauline R. Foncannon, and David H. Buzzell are with the Charleston Laboratory, Southeast Fisheries Center, National Marine Fisheries Service, NOAA, P.O. Box 12607, Charleston, SC 29412.

¹Relocated in 1978, this is now the Charleston Laboratory, Southeast Fisheries Center, NMFS, NOAA, P.O. Box 12607, Charleston, SC 29412.

Table 1 lists the values for the fat-soluble vitamins A, D, and E, and the water soluble vitamins choline, ascorbic acid, inositol, and biotin. The amount of vitamin A in the flesh is associated with the amount of fat present in the flesh. Dark flesh is higher in fat than the light; therefore, there is more vitamin A in the dark meat. Vitamin A is not synthesized by the aquatic animal, but is derived from the carotene in its foods. If the whole mollusk is used in the analyses, the amount of vitamin A will be influenced by the plankton in the intestinal tract of the animal.

So little data are reported on the amount of vitamin E present in the flesh of finfish or shellfish that no conclusion can be made. There is little evidence that more vitamin E is present in the flesh of the fatty fish.

The information on the three B-vitamins—choline, inositol, and biotin—is minimal. As noted in Table 1, raw fish flesh does contain varying amounts of vitamin C. In some fish, the amounts are large enough that one serving (100 g or 3.5 ounce) will make a sizable contribution to a person's daily requirement, ranging from 30 mg to 100 mg depending upon a person's body size.

Of the B-vitamins listed in Table 2, riboflavin and niacin are present in significant amounts; therefore, a serving of fish will contribute considerably to a person's daily needs. Thiamin content is found in notably lesser amounts. Mollusks, especially oysters, are rich in Vitamin B₁₂.

The data summarized in this report will provide the medical community with an estimate of the various vitamins found in seafoods for the calculations of special diets. The information will be also appreciated by those concerned with consumer education.

LITERATURE CITED

- Ackman, R. G., and M. G. Cormier. α -tocopherol in some Atlantic fish and shellfish with particular reference to live-holding without food. J. Fish. Res. Board Can. 24:357-373.
- Adrian, J., and R. Jacquot. 1956. Comparison between dried fish of Angola and industrial fish flours: Global composition, amino acids and B-complex vitamins (Title only.) Inter-African Nutr. Conf., Session III, p. 37.
- Ang, C. Y. W. 1974. Nutritional composition of frozen fish per 100 g. edible portion. Food Sci. Associates, Inc., Dobbs Ferry, N.Y.
- Arroba Nino, N. 1954. Analisis quimico bromatologico de la carne del *Neptomenus crassus* o "Cojinova" y de la *Sciaena deliciosa* o "Lorna". An. Fac. Farm. Bioquim. 5:115-117.
- Aschehoug, V., H. Kringstad, and G. Lunde. 1939. The vitamin D potency of different fish and fish products. J. Soc. Chem. Ind. 58:220-223.
- Bai, P. A., and M. Kalyani. 1960. Studies on ascorbic acid in fish tissues. I. J. Anim. Morphol. Physiol. 7(2):162-165.
- Bailey, B. E. 1943. Nutritive values of fishery products. Fish. Res. Board Can. Prog. Rep. Pac. Coast Stn. 57, p. 11.
- Bailey, B. E. (editor). 1952. Vitamins and other non-fat components of marine oils. Fish. Res. Board Can. Bull. 89:46-97.
- Bailey, B. E. 1952. Chart of the nutritive values of British Columbia fishery products. West. Fish. 6:50-54.
- Banerjee, D. K., and J. B. Chatterjee. 1963. Vitamin B₁₂ content of some articles of Indian diets and effect of cooking on it. Br. J. Nutr. 17:385-389.
- Banerjee, D. K., and J. B. Chatterjee. 1963. Effect of cooking on the availability of folic acid in Indian fishes. Bull. Calcutta School Trop. Med. 11(2):49-50.
- Banerjee, D. K., and J. B. Chatterjee. 1963. Vitamin B₁₂ content of Indian fishes and the effect of boiling on its availability. Bull. Calcutta School Trop. Med. 11(1):11-12.
- Barrett, I., and A. R. Connor. 1962. Blood lactate in yellowfin tuna, *Neothunnus macropterus*, and skipjack, *Katsuwonus pelamis*, following capture and tagging. Inter-Am. Trop. Tuna Comm. Bull. 6:231-280.
- Berezin, N. T. 1967. Use of fish and sea products in nutrition. Pishch. Promysh. Moskva, SFCSI-Int. TT-69-55095, p. 11-12.
- Booher, L. E., E. R. Hartzler, and E. M. Hewston. 1942. A compilation of the vitamin values of foods in relation to processing and other variants. U.S. Dep. Agric., Circ. 638, 244 p.
- Braekkan, O. R. 1965. Vitaminer i Norsk fish. IV. Vitamin B₆ og biotin forskjellige organer fra torskfisher (Gadidae) fanget langs Norskekysten. Fiskeridir. Skr. Ser. Teknol. Unders. 4(12), 11 p.
- Braekkan, O. R. 1956. Function of the red muscle in fish. Nature (Lond.) 178:747-748.
- Braekkan, O. R. 1958. Vitamin B₁₂ in marine fish. Nature (Lond.) 182:1386.
- Braekkan, O. R. 1962. B-vitamins in fish and shellfish. In E. Heen and R. Kreuzer (editors), Fish in nutrition, p. 132-139. Fishing News (Books) Ltd., Lond.
- Brooke, R. O., E. M. Ravesi, D. F. Gadbois, and M. A. Steinberg. 1966. Preservation of fresh unfrozen fishery products by low-level radiation. 5. The effects of radiation pasteurization on amino acids and vitamins in haddock filets. Food Technol. 20:1479-1482.
- Brooke, R. O., E. M. Ravesi, D. F. Gadbois, and M. A. Steinberg. 1964. Preservation of fresh unfrozen fishery products by low-level radiation. III. The effects of radiation on pasteurization on amino acids and vitamins in clams. Food Technol. 18(7):116-120.
- Brooks, G., and R. Paulais. 1939. Répartition et localisation des caroténoïdes, des flavines et de l'acid-l'ascorbique chez les mollusques lamellibranches: cas des huîtres et des gryphées vertes et blanches. C. R. Acad. Sci. 208:833-835.
- Butler, C. 1958. Nutritional value of fish in reference to atherosclerosis and current dietary research. Commer. Fish. Rev. 20(7):7-16.
- Cheftel, H. 1959. La valeur alimentaire des conserves appertisées. Etablissement J. J. Carnoud et Forges de Basse-Indie, Paris, Bull. 11.
- Cheldelin, V. H., and R. R. Williams. 1943. Studies of the average American diet. II. Riboflavin, nicotinic acid and pantothenic acid content. J. Nutr. 26:417-430.
- Cheldelin, V. H., A. M. Woods, and R. J. Williams. 1943. Losses of B vitamins due to cooking of foods. J. Nutr. 26:477-485.
- Collazos Chiriboga, C. 1956. La composición de los alimentos Peruanos. Arch. Venez. Nutr. 8(1):129-166.
- Dann, W. J., and P. Handler. 1942. The nicotinic acid content of meat. J. Nutr. 24:153-158.
- De Manes Braganaca, B. 1944. Nicotinic acid content of fish. Ann. Biochem. Exp. Med. (Calcutta) 4(1):41-44.
- Edisbury, J. R., J. A. Lovern, and R. A. Morton. 1937. LVIII. Distribution of vitamin A in the tissues of the eels *Anguilla vulgaris* and *A. aucklandii* Rich. Biochem. J. 31:416-423.
- Edwards, L. E., R. R. Sealock, W. W. O'Donnell, G. R. Bartlett, M. B. Barclay, R. Tully, R. H. Tybout, J. Box, and J. R. Murlin. 1946. Biological value of proteins in relation to the essential amino acids which they contain. IV. The analysis of fifteen protein foods for the ten essentials. J. Nutr. 32:597-612.
- Engel, R. W. 1943. The choline content of animal and plant products. J. Nutr. 25:441-446.
- Eusebio, E. C., and J. G. Palad. 1966. Vitamin A content of Philippine foods. Philipp. J. Sci. 95:199-207.
- Fieger, E. A. 1956. Vitamin content of fresh, frozen oysters. Quick Frozen Foods 19(4):152-155.
- Ferreira, F. A. G. 1951. Composition and nutritive value of some species of fish. Bol. Pesc. (Lisbon) 8:89-102.
- Fixsen, M. A. B., and M. H. Roscoe. 1938. Tables of the vitamin content of human and animal foods. Nutr. Abstr. Rev. 7:823-867.
- Fraga, F. 1958. Variación estacional de la composición química del mejillón (*Mytilus*

Continued on page 14.

Table 1.—Composition of the edible portion of raw (fresh or frozen) crustaceans, finfish, and mollusks. IV. Vitamins: A, D, E, choline, ascorbic acid, inositol, and biotin.

	A (I.U.)	D (I.U.)	E (mg)	Choline (mg)	Ascorbic acid (mg)	Inositol (μ g)	Biotin (μ g)	References
	per 100 g							
Abalone	61							64
Haliotidae spp.	1							
Amberjack and yellowtail	11 ± 11	27 ± 27	0.2 ± 0	59	2.0 ± 0.4			48, 55, 61, 62, 80, 103
Seriola spp.	$20-22$ 32	0-53 2	0.1-0.2 5	1	1.0-2.8 4			
Anchovies	635							33
Engraulidae spp.	1							
Barracudas	51 ± 7							57
Sphyrnidae spp.	40-62 3							
Basses, sea	129 ± 36							33, 57
Serranidae spp.	61-184 3							
Breams and porgies	53 ± 31			31 ± 4	2.9 ± 0.5			33, 57, 61, 80, 82
Sparidae spp.	7-144 4			18-39 3	0.4-1.7 4			
Butterfishes	182							69
Stromateidae spp.	1							
Carp	262 ± 127		630	2.0 ± 0.4	18.5 ± 6.9	$8,894 \pm 823$		36, 55, 57, 61, 62,
Cyprinidae spp.	29-1,020 8		1	0.6-3.1 6	0-160 24	7,585-11,000 4		67, 81, 83, 87, 122, 130, 140
Catfishes, airbreathing	633							33
Clariidae spp.	1							
Catfishes, freshwater		500			9.3 ± 1.9	6,620		36, 122, 130
Ictaluridae spp.		1			6.0-12.5 4	1		
Catfishes, sea	96				7.8 ± 3.9			33, 36, 83, 122
Ariidae spp.	1				0-11.7 3			
Characins					0 \pm 0			27
Characidae spp.					0-0 2			
Cichlids	58 ± 41				1.0			6, 33, 80
Cichlidae spp.	17-99 2				1			
Clams, mactra	33							80
Macridae spp.	1							
Clams, razor	64 ± 47							80, 81
Solenidae spp.	17-110 2							
Clams, Veneridae	$1,124 \pm 237$				14.2 ± 1.1			33, 44, 114
Veneridae spp.	887-1,360 2				13.1-15.2 2			
Clams	370 ± 172	4.4 ± 4.4	0.5 ± 0.2	58 ± 0	11.2 ± 3.6		2.3	1, 7, 33, 36, 40, 64,
Miscellaneous species	11-1,317 7	0-8.8 2	0-0.9 4	58-58 2	2.0-30.0 7		1	78, 80, 86, 101, 103
Codfish	25		0.2 ± 0				1.2 ± 0.7	1, 16, 19, 67, 86
Gadidae spp.	1		0.2-0.2 2				0.2-2.6 3	
Crabs	$5,115 \pm 3,755$				7.7 ± 5.4			33, 44, 85, 114
Mixed species	1,360-8,870 2				2.2-13.1 2			
Cusk							2.5 ± 2.0	16, 19
Brosme brosme							0.5-4.5 2	
Cusk eels and							0.1	141
broctulas								
Ophidiidae spp.							1	

Table 1.—Continued.

	A (I.U.)	D (I.U.)	E (mg)	Choline (mg)	Ascorbic acid (mg)	Inositol (μ g)	Biotin (μ g)	References
	per 100 g							
Cuttlefishes					2.3 \pm 0.7			42, 79, 80, 97,
Miscellaneous species					0.9-5.3			113, 147
					6			
Dace					1.0			87
Cyprinidae spp.					1			
Dolphins	179 \pm 154							55, 80
Coryphaenidae spp.	26-333							
	2							
Drums	199 \pm 155				10.2 \pm 3.9			4, 36, 57, 69,
Sciaenidae spp.	31-508				2.5-14.1			85, 122
	3				3			
Eels, conger					1.0			42
Congridae spp.					1			
Eels, freshwater	3,295 \pm 764	200			1.0 \pm 0.8			8, 30, 36, 61,
Anguillidae spp.	1,500-5,700				0.2-1.7			67, 80, 81, 96
	6	1			2			
Eels, snake	17							80
Ophichthidae spp.								
	1							
Eels, swamp	62 \pm 19							33, 80
Flutidae spp.	43-81							
	2							
Flounders, lefteye	335 \pm 182	43	140		1.0	7,519 \pm 609		48, 57, 62, 80, 130
Bothidae spp.	114-695					6,015-8,520		
	3	1	1		1	4		
Flounders, righteye	48 \pm 13	60	0.4		1.0 \pm 1.0		1.2	1, 3, 48, 80, 83, 111
Pleuronectidae spp.	34-73				0.0-2.0			
	3	1	1		2		1	
Goatfishes	104							33
Mullidae spp.								
	1							
Gobies	891				0.5 \pm 0			6, 33, 42
Gobiidae spp.					0.5-0.5			
	1				2			
Greenlings	0							7
Hexagrammidae spp.								
	1							
Grunts	136							57
Pomadasyidae spp.								
	1							
Gurnards, flying	59				0.0			27, 57
Dactylopteridae								
	1				1			
Haddock	50				0.0		2.6 \pm 2.3	13, 19, 86, 138
<i>Melanogrammus aeglefinus</i>							0.2-4.8	
	1				1		2	
Hake					1.0			35
Gadidae spp.								
					1			
Halibut	440	44			0.0	8,350 \pm 5,407	8.1 \pm 1.5	9, 26, 83, 130
Pleuronectidae spp.						1,400-19,000	6.6-9.5	
	1	1			1	3	2	
Herrings	814 \pm 620	1,627			9.0 \pm 3.8	9,590		8, 35, 69, 78,
Clupeidae spp.	100-4,531				0.0-27.7			81, 83, 88, 89,
	7	1			9	1		95, 122, 130
Jacks	125 \pm 53				3.3			33, 61
Carangidae spp.	72-177							
	2				1			
Lampreys	30,060 \pm 6,584	260 \pm 140						36, 51, 148
Petromyzontidae spp.	9,060-44,300	120-400						
	6	2						

Table 1.—Continued.

	A (I.U.)	D (I.U.)	E (mg)	Choline (mg)	Ascorbic acid (mg)	Inositol (μ g)	Biotin (μ g)	References
----- per 100 g -----								
Lings							1.1 \pm 0.1	16
<i>Molva</i> spp.							1.0-1.2 2	
Lizardfishes					3.0			80
Synodontidae spp.					1			
Lobsters and crayfishes	17				3.0 \pm 0.9		5.0 \pm 0.2	42, 78, 80, 83,
Mixed species	1				0.0-5.0 6		4.8-5.2 2	86, 87
Mackerels	107 \pm 40	1,036 \pm 273		11.1 \pm 4.4	3.0	8,060		5, 8, 15, 33, 48,
Scombridae spp.	0-711 17	143-2,000 6		2.5-29.2 6	1	1		57, 58, 67, 80, 81, 89, 95, 130
Milkfishes	537 \pm 137							33, 80
Chanidae spp.	400-673 2							
Minnows	79							57
Cyprinidae spp.	1							
Mojarras	62							33
Gerreidae spp.	1							
Mussel	1,226 \pm 1,193		0.5 \pm 0		4.1 \pm 1.4			15, 33, 36,
Mytilidae spp.	33-2,418 2		0.4-0.6 3		1.1-9.0 6			42, 53, 80
Octopuses					0			83
Mixed species					1			
Oysters	273 \pm 36	40			10.7 \pm 5.4		41 \pm 31	33, 42, 46, 78, 80,
Ostreidae spp.	170-366 5	1			0-38.1 7		10-72 2	81, 83, 86, 95, 145
Perches					1.2 \pm 0.0			87
Percidae spp.					1.1-1.2 2			
Perches, climbing	218							80
Anabantidae spp.	1							
Plaice							90	19
Pleuronectidae spp.							1	
Pollock	32						3.7 \pm 2.4	3, 16
Gadidae spp.	1						1.3-6.1 2	
Pompanos	5.1				0.0			4, 27
Carangidae spp.	1				1			
Sardines and pilchards	156 \pm 51	2,310 \pm 873						15, 33, 48,
Clupeidae spp.	50-303 5	532-5,400 5						103, 140
Sauries	1,551 \pm 1,493				2.0			57, 80
Scomberesocidae spp.	58-3,044 2				1			
Sawfish					0.0			83
Pristidae spp.					1			
Scad and mackerels	60 \pm 33		0.4		1.4 \pm 0.6			33, 35, 48,
Carangidae spp.	11-122 3		1		0.8-2.0 2			57, 62, 80
Scallops	0				3.0		0.3	78, 86
Pectinidae spp.	1				1		1	
Sea chubs	19							57
Kyphosidae spp.	1							

Table 1.—Continued.

	A (I.U.)	D (I.U.)	E (mg)	Choline (mg)	Ascorbic acid (mg)	Inositol (μ g)	Biotin (μ g)	References
	per 100 g							
Sea cucumbers					0.0			40
Mixed species					1			
Sea robins					1.0			42
Triglidae spp.					1			
Shad	138							33
Clupeidae spp.	1							
Sharks, dogfish	797 \pm 211	15						8, 49
Squalidae spp.	114-1,600							
	8	1						
Sharks			270.0		0.0			27, 62
Miscellaneous species			1		1			
Sheathfishes	67							80
Siluridae spp.	1							
Shrimp and prawns	108 \pm 35	150 \pm 0			1.5 \pm 0.6	542 \pm 319	1.0	5, 15, 27, 33, 36, 64,
Mixed species	20-297	150-150			0.0-3.0	4-2,400		76, 78, 80, 81, 83, 86, 98
	8	2			5	7	1	
Silversides					1.0			83
Atherinidae spp.					1			
Smelts	96				1.4 \pm 1.4			57, 61
Osmeridae spp.	1				0.0-2.8			
					2			
Snails	83				15.0			80, 118
Mixed species	1				1			
Snappers	187				0.3			65, 69
Lutjanidae spp.	1				1			
Snooks					0.0			83
Centropomidae spp.					1			
Soapies	33							80
Leiognathidae spp.	1							
Soles	39							3
Pleuronectidae spp.	1							
Spadefishes	98							33
Ephippidae spp.	1							
Squid					4.9			113
Mixed species					1			
Surfperches	62 \pm 17							57
Embiotocidae spp.	45-79							
	2							
Surgeonfishes					1.3			61
Acanthuridae spp.					1			
Therapons	29							57
Theraponidae spp.	1							
Threadfins	172			8.0 \pm 0.0				10, 12, 69
Polynemidae spp.	1			8.0-8.0				
				2				
Tilefishes	17					6,350		80, 130
Branchiostegidae spp.	1					1		

Table 1.—Continued.

	A (I.U.)	D (I.U.)	E (mg)	Choline (mg)	Ascorbic acid (mg)	Inositol (μg)	Biotin (μg)	References
	per 100 g							
Trout, salmon Salmonidae spp.	173±49 3-600 17	360±107 30-1,200 12			7.1±1.9 1.3-9.0 4	19 1	3.7±1.4 0.9-5.3 3	7, 8, 9, 15, 19, 26, 46, 53, 55, 78, 81, 86, 89, 95, 120
Trout, trout Salmonidae spp.	41±28 13-69 2			87 1	1.1±1.1 0.0-2.2 2			31, 53, 83, 111
Tuna Scombridae spp.	129±66 0-963 14	1,125±307 700-2,000 4		37 1	2.6±1.2 0.0-10.7 9		1.5 1	8, 15, 27, 33, 39, 46, 57, 69, 78, 80, 83, 86, 89, 95, 102
Turbot Turbanidae spp.	233 1				2.0 1			80
Turbots Pleuronectidae spp.	39 1							3
Whales <i>Physeter macrocephalus</i>	267 1							15
Whitings Sillaginidae spp.	54±54 0-107 2							33, 36
Wrasses Labridae spp.	48 1							57

¹Average of the means.²Range of the means.³Number of mean values.Table 2.—Composition of the edible portion of raw (fresh or frozen) crustaceans, finfish, and mollusks. IV. Vitamins: Thiamin, riboflavin, niacin, pyridoxine, folic acid, B₁₂, and pantothenic acid.

	Thiamin (μg)	Riboflavin (μg)	Niacin (mg)	Pyridoxine (μg)	Folic acid (μg)	B ₁₂ (μg)	Pantothenic acid (μg)	References
	per 100 g							
Abalone Haliotidae spp.	1210±30 180-240 2	85±27 56-140 3	1.4±0.1 1.2-1.6 3	0.6 1		0.7 1	2,300±0 2,300-2,300 2	50, 64, 80, 81, 100, 151
Amberjacks and yellowtails <i>Seriola</i> spp.	115±33 27-180 4	84±17 40-120 4	8.3±8 6.8-10 4		3.5±0.3 3.2-3.7 2	4.2±1.4 1.3-5.6 3	598±3 595-600 2	2, 50, 64, 80, 85, 102
Anchovies Engraulidae spp.	19±7 8-37 4	104±24 79-152 3	3.1±0.4 2.3-3.5 3	181±27 144-260 4	12.3 1	6.3±0 6.3-6.3 2		44, 64, 66, 80, 84, 85, 93, 111, 116, 134, 153
Augar shells <i>Terebralia aulacata</i>					21.1 1	12.6 1	60 1	80
Barracudas Sphyraenidae spp.	92±21 44-134 4	70±11 27-90 5	4.1±0.6 2.4-5.9 5	350±101 150-470 3	11.9 1	1.4±0.4 0.2-1.8 4		64, 80, 82, 85, 88, 150, 153, 154
Basses, sea Serranidae spp.	79±13 20-170 14	114±33 35-374 13	2.4±0.4 1.1-4.3 12	300 1	8.8 1	5.0±2.4 0.3-9.1 4	7,530 1	3, 19, 27, 43, 44, 64, 65, 80, 82, 85, 130
Basses, temperate Percichthyidae spp.	100 1	52 1	2.4 1					44
Bigeyes Scombroptidae spp.				400 1				155
Billfishes Istiophoridae spp.	100 1	60 1	4.5 1					80

Table 2.—Continued.

	Thiamin (μ g)	Riboflavin (μ g)	Niacin (mg)	Pyridoxine (μ g)	Folic acid (μ g)	B ₁₂ (μ g)	Pantothenic acid (μ g)	References
	per 100 g							
Bluefishes Pomatomidae spp.	135 \pm 15 120-150 2	120 \pm 30 90-150 2	1.9 \pm 0 1.9-1.9 2					23, 82, 146
Bombay duck <i>Harpodon nohereus</i>	30 \pm 10 20-40 2	70 \pm 20 50-90 2	5.7 \pm 5.1 0.6-10.7 2	970 1				66, 80
Breams and porgies Sparidae spp.	77 \pm 18 10-190 15	96 \pm 21 20-250 15	5.4 \pm 6 1.5-7.8 12	460 \pm 0 460-460 2	1.6 \pm 1.2 0.4-4 3	1.8 \pm 0.4 0.1-2.8 7	213 \pm 8 205-220 2	2, 50, 64, 66, 85, 153
Burbot Gadidae spp.	388 \pm 27 306-455 6	141 \pm 1 140-142 2	1.6 \pm 0 1.6-1.6 2					43, 63, 71, 72, 139
Butterfishes Stromateidae spp.	163 \pm 1 163-164 2	223 \pm 109 90-550 4	4.8 \pm 1.2 2.6-8.1 4	450 1		2.3 \pm 0 2.3-2.3 2		29, 70, 80, 85, 116, 126
Butterflyfishes Chaetodontidae spp.	33 \pm 4 25-40 3	45 \pm 26 19-70 2	4.5 \pm 0.1 4.4-4.6 3					44, 64, 80, 109
Caesios Caesionidae spp.	24 \pm 4 20-28 2	24 \pm 4 20-27 2	3.8 \pm 0.6 3.2-4.3 2	190 1		2.3 \pm 0.8 1.5-3.1 2		50, 64, 80, 85
Carp Cyprinidae spp.	114 \pm 32 5-448 20	81 \pm 23 10-180 7	3.0 \pm 0.7 1.3-11 16	71 \pm 60 11-130 2	70 \pm 68 2.8-138 2	3.2 \pm 1.2 0.1-9 8	150 1	10, 11, 12, 29, 63, 71, 72, 80, 85, 109, 110, 111, 134, 136, 137, 149, 153, 154
Catfishes, airbreathing Clariidae spp.	44 \pm 23 8-110 4	36 \pm 5 31-40 2	1.7 \pm 0.6 0.5-3.2 4		13.6 \pm 1.7 12.0-15.3 2	3.5 \pm 0.1 3.4-3.7 4	460 1	10, 11, 29, 36, 64, 80, 85, 109
Catfishes, freshwater Ictaluridae spp.	0 1		1.8 \pm 0.5 1.0-2.5 3		3.1 \pm 1.2 1.9-4.3 2	3.7 \pm 0.4 2.2-4.6 7	464 \pm 4 460-468 2	10, 11, 29, 50, 111, 116, 124, 135
Catfishes, sea Ariidae spp.	61 \pm 8 40-80 4	115 \pm 22 80-197 5	2.4 \pm 0.6 0.5-4.5 6	370 1	82.5 \pm 67.5 15-150 2	2.4 \pm 0.2 2.2-2.5 2	570 1	64, 109, 124, 138
Characins Characidae spp.	30 \pm 12 10-50 3	87 \pm 19 50-110 3	3.7 \pm 1.4 1.7-6.3 3			1.6 1		27, 65
Chimaeras Chimaeridae spp.					2.7 1		310 1	50
Cichlids Cichlidae spp.	30 \pm 0 30-30 2	85 \pm 35 50-120 2	2.3 \pm 0.8 1.5-3.1 2	320 1		2.9 \pm 0.2 2.7-3.0 2		82, 85
Clams, arkshells Arcidae spp.	2 1	200 1	1.5 1		17 1	13.6 \pm 6.4 7.2-20 2	790 1	50, 64, 151
Clams, freshwater Corbiculidae spp.					17 1	12.1 1	360 1	33, 50, 151
Clams, maestra Mactridae spp.	72 \pm 52 20-128 2	114 \pm 16 98-130 2				0.4 \pm 0.2 0.2-0.7 3		25, 151
Clams, razor Solenidae spp.	95 \pm 5 90-100 2	355 \pm 155 180-490 2	1.6 \pm 0.1 1.5-1.6 2			8.6 1		80, 81, 151
Clams, softshell Myacidae spp.	21 \pm 19 1-79 4	180 \pm 29 95-219 4	1.9 \pm 0.4 1.2-2.5 3	88 \pm 9 77-106 3		71.5 \pm 22.4 7.1-105.0 4	293 \pm 17 260-311 3	21, 59, 121, 125
Clams, Tellin Tellinidae spp.	10 1	20 1	2.2 1	70 1	10.9 1			80
Clams Veneridae spp.	100 \pm 33 10-240 7	381 \pm 147 20-940 7	2.9 \pm 0.6 1.1-5.0 6			3.9 \pm 1.7 2.2-5.6 2		23, 44, 64, 85, 109, 125

Table 2.—Continued.

	Thiamin (μ g)	Riboflavin (μ g)	Niacin (mg)	Pyridoxine (μ g)	Folic acid (μ g)	B ₁₂ (μ g)	Pantothenic acid (μ g)	References
	per 100 g							
Clams	49 \pm 12	238 \pm 47	1.3 \pm 0.2	75 \pm 42	25.1 \pm 6.5	9.8 \pm 2.5	531 \pm 91	23, 45, 64, 75, 78, 80,
Miscellaneous species	2-139 14	12-780 16	0.2-2.3 9	0-350 8	2.7-58.0 7	0.2-62.3 34	440-622 2	86, 94, 100, 111, 123, 125, 127, 142, 151
Codfishes	71 \pm 9	121 \pm 23	2.5 \pm 0.4	221 \pm 25	5.0 \pm 1.6	0.6 \pm 0.2	163 \pm 20	14, 16, 18, 19, 23, 36,
Gadidae spp.	18-150 17	11-325 16	0.2-6.7 14	170-288 4	1.8-6.7 3	0.1-2 11	96-400 14	43, 50, 59, 60, 73, 80, 81, 86, 90, 97, 110, 111, 112, 125, 134, 138, 141, 144, 146, 155
Cornet						0		153
Fistulariidae spp.						1		
Crabs	85 \pm 20	382 \pm 280	2.5 \pm 0.2					23, 43, 44, 90
Mixed species	10-140 6	60-940 3	2.3-2.7 2					
Croakers	85 \pm 26	98 \pm 19	3.2 \pm 0.9	0.2		2.5		80, 81, 92
Sciaenidae spp.	40-160 4	60-150 4	1.7-5.5 4	1		1		
Cusk	38 \pm 6	303 \pm 160	2.7 \pm 0.2	304 \pm 30		1.0	261 \pm 44	16, 19, 23, 90, 138, 146
<i>Brosme brosme</i>	32-51 3	94-940 5	2.3-3.0 3	274-333 2		1	273-310 3	
Cusk eels and brotulas	175	35 \pm 3	1.8 \pm 0.4		2.0	0.6 \pm 0.3	115	65, 141
Ophidiidae spp.	1	32-37 2	1.4-2.2 2		1	0.3-0.9 2	1	
Cuttlefishes	96 \pm 59	480 \pm 430	2.1 \pm 0.8					70, 80, 113, 147
Miscellaneous species	9-210 3	50-910 2	1.2-2.9 2					
Dolphins	20	70	6.1			0.1		80, 153
Coryphaenidae spp.	1	1	1			1		
Dories				952		0		93, 153
Zeidae spp.				1		1		
Drepanes	29 \pm 2	111 \pm 30	3.2 \pm 0.4					64, 80
Drepanidae spp.	27-30 2	81-140 2	2.8-3.7 2					
Drums	63 \pm 14	131 \pm 46	3.1 \pm 0.7			2.4 \pm 1.5		2, 27, 59, 64, 65,
Sciaenidae spp.	20-130 9	31-530 10	0.5-8.9 11			0.2-5.3 3		70, 102, 124
Eels, arrowtooth				230			150	127
Dysommidae spp.				1			1	
Eels, conger	60	40 \pm 0	4.3 \pm 0.8			3	240	80, 138
Congridae spp.	1	40-40 2	3.5-5.0 2			1	1	
Eels, freshwater	191 \pm 31	335 \pm 51	2.3 \pm 0.5	254 \pm 15	11.6 \pm 1.6	1.0 \pm 0	141 \pm 6	19, 50, 59, 61, 80, 81,
Anguillidae spp.	144-280 4	190-520 6	1.4-3.5 4	230-300 5	10.0-13.1 2	1.0-1.0 3	125-150 4	84, 96, 111, 119, 134
Eels, moray	10	30	3.1			0.3 \pm 0.2		64, 153
Muraenidae spp.	1	1	1			0.1-0.4 2		
Eels, pike conger	60	90	2.7			0.2		80, 153
Muraenesocidae spp.	1	1	1			1		
Eels, snake	41 \pm 7	120	1.3 \pm 0.5			1.5 \pm 0.5		10, 12, 80, 85,
Ophichthidae spp.	30-53 3	1	0.5-2.3 3			1.0-2.6 3		109, 124, 149
Eels, spiny	60							109
Motacanthidae spp.	1							
Eels, swamp	85 \pm 45	135 \pm 85	2.2 \pm 0.4					33, 80
Flutidae spp.	40-130 2	50-220 2	1.8-2.5 2					
Featherbacks	85 \pm 35	55 \pm 15	3.3 \pm 2.6			3.4		29, 80, 135, 136, 137
<i>Notopterus</i> spp.	50-120 2	40-70 2	0.4-8.5 3			1		

Table 2.—Continued.

	Thiamin (μ g)	Riboflavin (μ g)	Niacin (mg)	Pyridoxine (μ g)	Folic acid (μ g)	B ₁₂ (μ g)	Pantothenic acid (μ g)	References
	per 100 g							
Flatheads Percophidae spp.	20 \pm 1 19-20 2	60 \pm 30 30-90 2	4.7 \pm 0.4 4.3-5.0 2				240	64, 80
Flounders, lefteye Bothidae spp.	85 \pm 15 70-100 2	73 \pm 27 46-100 2	5.3 \pm 0.4 5.0-5.7 2	130 1		1.3 1	900 1	65, 80, 153, 154
Flounders, righteye Pleuronectidae spp.	103 \pm 31 30-400 11	84 \pm 23 40-335 12	2.3 \pm 0.4 0.8-3.8 9	191 \pm 24 137-250 4	5 \pm 0 5-5 2	5.4 \pm 4.6 0.8-10 2	863 \pm 240 245-1,700 6	3, 19, 50, 59, 80, 81, 83, 84, 97, 110, 111, 115, 125, 134, 138, 146
Flyingfishes Exocoetidae spp.	17 \pm 7 4-26 3	56 \pm 10 40-80 4	3.9 \pm 0.6 3.0-5.6 4		2.8 1	1.0 \pm 0.4 0.3-1.3 3		64, 80, 85, 153
Gars Lepisosteidae spp.				900 1				134
Goatfishes Muliidae spp.	36 \pm 6 25-43 3	59 \pm 21 38-80 2	1.5 \pm 0.1 1.4-1.6 2	1.7 1				64, 80, 93, 109
Greenlings Hexagrammidae spp.	72 \pm 15 43-105 4	49 \pm 8 38-73 4		540 1	7.4 \pm 0 7.4-7.4 2	13.2 \pm 4.8 3.6-18.0 3	188 \pm 3 185-190 2	50, 67, 80, 111, 125, 140
Grunts Pomadasysidae spp.	77 \pm 23 18-200 7	226 \pm 126 32-900 7	4.2 \pm 0.9 2.0-8.6 7	200 1	7.0 \pm 3.5 3.5-10.5 2	0.4 \pm 0.3 0.1-0.7 2	270 \pm 0 270-270 2	27, 44, 50, 64, 65, 66, 80, 153, 154
Gumards, flying Dactylopteridae spp.	85 \pm 75 10-160 2	75 \pm 5 70-80 2	2.8 \pm 0.7 2.1-3.4 2	480 1	1.8 1	0.3 1	225 1	27, 50, 80, 153, 154
Haddock <i>Melanogrammus aeglefinus</i>	39 \pm 5 9-100 19	69 \pm 11 12-210 24	3.6 \pm 0.1 2.4-4.3 20	231 \pm 16 122-300 11	0.8 1	1.3 \pm 0.1 0.5-3.5 19	145 \pm 17 49-380 20	18, 19, 20, 23, 43, 59, 60, 83, 86, 90, 111, 119, 127, 138, 141, 147
Hake Gadidae spp.	89 \pm 27 39-132 3	75 \pm 5 70-80 2	1.6 \pm 0.6 1.0-2.2 2	875 1				35, 44, 93, 119
Halibut Pleuronectidae spp.	83 \pm 11 40-180 13	80 \pm 11 44-185 15	7.5 \pm 1.0 2.8-14.2 11	400 \pm 15 347-430 5	2.6 \pm 0.3 2.0-2.9 3	0.8 \pm 0.1 0.7-1.0 6	303 \pm 50 111-595 11	7, 9, 19, 23, 25, 26, 50, 59, 77, 80, 83, 84, 86, 90, 110, 111, 125, 130, 134, 138, 141
Herring Clupeidae spp.	46 \pm 8 6-170 27	261 \pm 39 50-1,000 26	3.8 \pm 0.4 0.6-9.6 25	310 \pm 43 160-450 6	10.3 \pm 2.9 1.7-14.0 4	11.4 \pm 4.3 1.4-34.0 9	2,427 \pm 1,415 970-9,500 6	2, 9, 10, 11, 12, 14, 27, 29, 31, 36, 59, 64, 66, 67, 70, 73, 78, 80, 81, 82, 84, 95, 97, 110, 111, 117, 124, 125, 127, 138, 153, 154
Jacks Carangidae spp.	59 \pm 17 15-122 5	60 \pm 10 25-81 5	5.7 \pm 0.5 3.9-7.1 5	670 1		7.5 \pm 1.6 5.9-9.1 2		64, 80, 85
Lampreys Petromyzontidae spp.	339 \pm 256 46-850 3	520 \pm 93 427-612 2	4.7 1	195 \pm 25 170-220 2	25.5 \pm 1.7 23.6-29.0 3	4.1 \pm 0.3 3.8-4.4 2	425 \pm 145 280-570 2	50, 51, 80
Lings <i>Molva</i> spp.		80 1	2.5 \pm 0.2 2.3-2.7 2	265 \pm 44 221-309 2		1 \pm 0.5 0.5-1.9 3	320 1	10, 18, 97, 138
Lizardfishes Synodontidae spp.	92 \pm 12 80-104 2	39 \pm 11 28-50 2	2.9 \pm 0.5 2.4-3.4 2					64, 80
Lobsters and crayfishes Miscellaneous species	99 \pm 24 7-165 7	64 \pm 10 10-130 10	2.3 \pm 0.3 1.2-4.3 8	210 1	0.6 \pm 0 0.6-0.6 2	1.6 \pm 0.6 0.5-2.7 4	410 \pm 0 410-410 2	38, 59, 64, 78, 80, 83, 86, 90, 111, 138, 141, 149
Longarays Ambasidae spp.	11 1	39 1	1.5 1					64

Table 2.—Continued.

	Thiamin (μ g)	Riboflavin (μ g)	Niacin (mg)	Pyridoxine (μ g)	Folic acid (μ g)	B ₁₂ (μ g)	Pantothenic acid (μ g)	References
	per 100 g							
Mackerels Scombridae spp.	90 \pm 11 10-237 31	263 \pm 49 27-940 30	12.4 \pm 2.0 2.7-58.0 27	654 \pm 76 270-1,000 10		4.2 \pm 1.2 0.4-16.2 14	444 \pm 82 160-850 8	17, 23, 38, 43, 44, 50, 52, 54, 59, 64, 65, 66, 74, 80, 81, 85, 90, 101, 102, 105, 107, 109, 111, 112, 127
Milkfishes Chanidae spp.	57 \pm 44 13-100 2	77 \pm 23 54-100 2	6.1 \pm 0.3 5.8-6.4 2	420 1	15.9 1	3.4 \pm 0 3.4-3.4 2		64, 80, 85
Minnows Cyprinidae spp.	30 1	100 1	3.5 1	130 1				80, 154
Mojarras Gerreidae spp.	51 \pm 21 30-92 3	85 \pm 3 80-90 3	5.0 \pm 0.3 4.5-5.3 3	360 1	21.4 1	2.4 \pm 0.5 1.9-3.4 3		64, 65, 80, 85
Mussels Mytilidae spp.		150 \pm 30 120-180 2	1.6 \pm 0.4 1.2-2.0 2	98 \pm 92 6-190 2	41.8 1	10.2 1		1, 42, 80, 85
Needlefishes Belonidae spp.	4 1	39 \pm 2 37-40 2	0.9 \pm 0 0.9-0.9 2	655 \pm 85 570-740 2		1.9 \pm 0 1.9-1.9 2		64, 80, 85, 111
Ocean perches Sebastes marinus		110 1	2.0 1				360 1	138
Octopuses Mixed species	67 \pm 26 20-140 5	73 \pm 14 40-110 5	3.2 \pm 1.2 1.3-5.3 3	360 1				80, 83, 111, 125
Oysters Ostreidae spp.	153 \pm 11 9-300 36	188 \pm 15 16-340 29	1.8 \pm 0.2 0.7-7.1 34	166 \pm 25 30-320 18	84.4 \pm 77.8 3.7-240 3	17.2 \pm 2.6 11.5-33.0 9	365 \pm 22 184-530 22	1, 9, 22, 23, 34, 38, 40, 45, 59, 64, 78, 79, 80, 81, 83, 85, 91, 94, 95, 97, 100, 111, 115, 121, 127, 141, 142, 145, 151
Parrotfishes Scaridae spp.	29 \pm 24 5-53 2	57 \pm 6 50-70 3	1.4 \pm 0 1.4-1.5 3	180 1		0.6 1		64, 65, 80
Perches Percidae spp.	187 \pm 63 60-250 3	119 \pm 46 28-170 3	1.9 \pm 0.2 1.4-2.3 4					19, 28, 30, 63
Perches, climbing Anabantidae spp.	39 \pm 12 19-99 6	274 \pm 49 190-361 3	2.4 \pm 0.7 0.8-3.7 4	200 1	2.7 \pm 1.3 1.0-5.2 3			11, 12, 36, 64, 80, 109, 124
Pikes Esocidae spp.				135 \pm 17 115-150 2				111, 134
Plaice Pleuronectidae spp.	137 \pm 32 105-200 3	163 \pm 33 130-195 2	4.0 1	217 \pm 14 182-250 4		1.4 \pm 0.4 1.0-2.2 3	800 \pm 0 800-800 2	19, 84, 90, 111, 134
Pollock Gadidae spp.	82 \pm 18 45-160 6	122 \pm 18 80-200 7	2.6 \pm 0.6 1.6-2.9 4	238 \pm 60 60-473 6	65 1	2.1 \pm 0.5 1.0-3.5 5	274 \pm 41 140-380 5	3, 16, 18, 23, 45, 50, 59, 67, 80, 90, 111, 138, 146, 154
Pomfrets Bramidae spp.	120 \pm 70 50-190 2	190 \pm 110 80-300 2	0.6 \pm 0.2 0.4-0.8 2					66, 80
Pompanos Carangidae spp.	254 \pm 154 90-562 3	91 \pm 17 60-118 3	4.7 \pm 1.5 3.0-8.1 3					27, 44
Rays, stingray Dasyatidae spp.	50 \pm 10 36-80 4	44 \pm 5 30-50 4	3.5 \pm 0.6 2.5-4.6 4			0.1 1		64, 80, 153
Rockfishes Scorpaenidae spp.	96 \pm 18 29-153 7	150 1	3.0 1	143 \pm 88 55-230 2	4.7 \pm 4.3 0.4-9 2	1.0 1	177 \pm 92 80-360 3	50, 80, 111, 125, 154
Sandfishes Trichodontidae spp.	100 1	50 1			7.8 \pm 0 7.8-7.8 2		630 \pm 0 630-630 2	50, 80

Table 2.—Continued.

	Thiamin (μ g)	Riboflavin (μ g)	Niacin (mg)	Pyridoxine (μ g)	Folic acid (μ g)	B ₁₂ (μ g)	Pantothenic acid (μ g)	References
	per 100 g							
Sardines and pilchards Clupeidae spp.	19-8 1-90 10	107-38 30-387 9	6.3-1.0 2.4-10 9	424-231 150-882 3	2 1	5.6-2.3 1.1-17.0 7	1,030-30 1,000-1,090 3	8, 15, 23, 33, 36, 45, 48, 50, 64, 80, 82, 85, 93, 103, 110, 111, 116, 140, 153, 154
Sauries Scomberesocidae spp.	50-1 48-54 5	112-11 87-132 5	6.0 1 1	660 1 1	6.4 1 1		850 1	50, 80, 99, 154
Sawfishes Pristidae spp.	140 1	190 1	4.1 1					83
Scad and mackerel Carangidae spp.	139-19 26-180 8	141-13 77-190 7	6.0-1.3 1.5-11.0 7	406-98 300-700 4	4.3-2.4 1.9-6.6 2	5.3-1.4 0.2-12.7 11	350-0 350-350 2	19, 45, 50, 57, 64, 80, 85, 101, 102, 103, 140, 149, 153, 154
Scallops Pectinidae spp.	85-45 40-130 2	83-18 65-100 2	1.3-0.1 1.2-1.4 2			8.9-7.6 1.2-24.0 3	138-6 132-143 2	78, 81, 94, 100, 111, 141
Scorpionfishes Scorpaenidae spp.				55 1		7.6-4.4 3.2-12.0 2	80 1	67, 111
Sculpins Cottidae spp.			3.2 1			6.1 1	100 1	50, 153
Sea cucumber Mixed species	53-18 35-70 2	90-0 90-90 2	0.5 1			1.4-0.3 0.5-2.0 4		45, 80, 94, 125
Sea robins Triglidae spp.	90 1							43
Shad Clupeidae spp.	46-35 10-150 4	240-0 240-240 2	8.4 1				608 1	23, 33, 59, 64, 80, 111
Shark, dogfish Squalidae spp.	52-8 40-66 3	264-155 80-573 3	2.9-1.2 1.0-5.2 3		3.2 1	1.8-0 1.8-1.8 2	747-57 690-860 3	43, 50, 70, 80, 90, 111, 138
Sharks, hammerhead Sphyrnidae spp.	14 1	81 1	3.2 1					64
Sharks, mackerel Lamnidae spp.	105-25 80-130 2	91-10 81-100 2	9.5-2.5 7.0-12.1 2			3.0-0.4 2.6-3.4 2	330 1	2, 138
Sharks, sand tiger Odontaspidae spp.	20 1	30 1	4.4 1					44
Sharks Miscellaneous species	40-16 5-110 6	57-15 19-110 6	4.0-0.9 0.9-6.6 6			0.4-0.2 0-0.7 3		27, 64, 65, 80, 102, 116
Sheathfishes Siluridae spp.	10 1	40 1	1.4 1					80
Shiners Cyprinidae spp.						1.4 1		10
Shrimp and prawns Mixed species	41-6 10-143 28	76-11 13-190 26	2.7-0.3 0.7-4.9 17	66-13 16-125 10	5.2-2.2 3.0-7.4 2	3.8-1.0 0.9-8.1 7	278-16 165-372 15	27, 36, 38, 64, 78, 80, 81, 83, 84, 85, 86, 90, 98, 108, 111, 125, 127, 134, 138, 141, 143, 149
Siganids Siganoidae spp.	203-43 160-245 2	129-2 127-130 2	4.8-0.4 4.4-5.1 2	140 1		0.1 1		64, 80, 153
Silversides Atherinidae spp.	88-71 10-230 3	65-15 50-80 2	2.9-1.6 1.3-4.5 2					43, 80, 83

Table 2.—Continued.

	Thiamin (μ g)	Riboflavin (μ g)	Niacin (mg)	Pyridoxine (μ g)	Folic acid (μ g)	B ₁₂ (μ g)	Pantothenic acid (μ g)	References
	per 100 g							
Sleepers Eleotridae spp.	26 1							109
Smelts Osmeridae spp.	54 \pm 26 10-130 4	143 \pm 33 43-360 8	1.9 \pm 0.4 1.3-3.0 4	120 1	5.0 \pm 1.3 3.7-6.3 2	1.8 \pm 1.6 0.2-3.4 2	919 \pm 281 638-1,200 2	15, 50, 59, 63, 80, 86, 125, 141, 153, 154
Snails Mixed species	56 \pm 54 2-110 2	169 \pm 111 58-280 2	1.5 \pm 0.2 1.3-1.6 2	120 1	31.8 1	26.7 \pm 7.1 9.0-77.0 10		64, 80, 85, 151
Snappers Lutjanidae spp.	88 \pm 18 46-170 6	81 \pm 17 38-131 5	3.6 \pm 0.6 2.5-5.2 5	100 \pm 100 0-200 2				43, 44, 64, 65, 80, 83
Snooks Centropomidae spp.	134 \pm 47 48-350 8	150 \pm 59 48-440 7	1.4 \pm 0.4 0.7-3.1 8		65.0 1	3.4 \pm 0 3.4-3.4 2		10, 11, 12, 29, 36, 44, 64, 66, 80, 83
Soapies Leiognathidae spp.	43 \pm 16 13-70 4	39.3 \pm 0.8 37-40 4	2.1 \pm 0.1 1.9-2.5 4	170 1	23.6 1	3.6 \pm 0 3.6-3.6 2		64, 80, 85
Soles Pleuronectidae spp.	62 \pm 5 38-88 10	48 \pm 2 37-57 7	1.1 1	1,045 1	3.0 \pm 0.1 2.9-3.1 2	0.1 1	420 \pm 197 155-805 3	3, 9, 50, 93, 111, 119, 125, 153
Soles Soleidae spp.	72 1							15
Spadefishes Ephippidae spp.	66 \pm 46 20-158 3	157 \pm 7 150-170 3	5.2 \pm 0.1 5.0-5.3 3			2.6 \pm 0 2.6-2.6 2		33, 64, 65, 80
Squid Mixed species	43 \pm 19 8-150 7	218 \pm 104 50-836 7	2.6 \pm 0.5 1.2-4.7 7	623 \pm 360 70-1,300 3	12.5 1	4.5 \pm 2.9 1.3-13.0 4	680 1	45, 54, 65, 80, 81, 93, 98, 100, 113
Suckers Catostomidae spp.	15 \pm 11 4-36 3	50 1	1.3 \pm 0.1 1.1-2.0 7				464 1	19, 44, 63, 71, 72, 111
Sunfishes Centrarchidae spp.			0.5 1				512 1	111, 124
Surgeonfishes Acanthuridae spp.	29 \pm 2 27-30 2	28 \pm 2 26-30 2	3.8 \pm 0 3.8-3.8 2		10 1	1.1 \pm 0.5 0.6-1.6 2	320 1	64, 80, 153
Tarpons Elopidae spp.	54 \pm 21 16-90 4	52 \pm 5 39-60 4	3.6 \pm 0.7 1.6-5.1 5					29, 64, 80
Therapons Theraponidae spp.	23 \pm 2.5 20-28 3	71 \pm 5 63-81 3		600 1		4.9 \pm 2.2 2.7-9.2 3		57, 64, 80, 85
Threadfins Polynemidae spp.	25 \pm 5 19-40 4	93 \pm 26 50-160 4	2.6 \pm 0.6 1.5-4.6 5		70 1			11, 64, 66, 79, 80
Tilefishes Branchiostegidae spp.	80 1	111 1	2.8 1	45 1				80, 154
Tonguefishes Cynoglossidae spp.	50 1	150 1	3.3 1					80
Trout, salmon Salmonidae spp.	130 \pm 15 30-348 23	143 \pm 13 46-231 26	7.2 \pm 0.2 5.6-8.8 15	745 \pm 54 590-975 6	3.9 \pm 0.4 2.2-4.8 6	1.7 \pm 0.5 0.1-5.0 13	988 \pm 144 490-2,080 15	15, 19, 23, 24, 25, 26, 43, 47, 53, 55, 59, 73, 78, 80, 81, 84, 95, 97, 101, 104, 106, 111, 120, 125, 127, 132, 134, 138, 141, 144, 146, 149, 153
Trout, trout Salmonidae spp.	80 \pm 9 10-140 12	109 \pm 17 10-210 15	4.2 \pm 0.6 2.5-7.8 9	690 \pm 0 690-690 2		3.3 \pm 0.8 1.0-5.0 6	1,843 \pm 77 1,630-2,000 5	19, 38, 44, 59, 61, 80, 83, 111, 125, 138, 152, 153

Table 2.—Continued.

	Thiamin (μ g)	Riboflavin (μ g)	Niacin (mg)	Pyridoxine (μ g)	Folic acid (μ g)	B ₁₂ (μ g)	Pantothenic acid (μ g)	References
	per 100 g							
Trout, whitefish Salmonidae spp.	88 \pm 5 27-126 26	120 \pm 0 120-120 3	2.8 \pm 0.2 0.6-4.6 15					19, 43, 63, 71, 72
Tunas Scombridae spp.	120 \pm 21 10-434 22	164 \pm 31 13-660 28	9.7 \pm 1.1 0-23.4 21	647 \pm 72 190-920 10	2.1 \pm 0.7 0.6-3.2 4	6.2 \pm 2.5 0.2-47.0 19	917 \pm 313 186-3,280 9	2, 15, 17, 18, 27, 45, 56, 64, 70, 74, 80, 82, 84, 89, 95, 105, 110, 127, 131, 134, 138, 141, 146, 147, 149, 153, 154
Turbons Turbanidae spp.	60 1	60 1	3.0 1	1,700 1	7.3 \pm 0 7.3-7.3 2	0.3 1	305 \pm 45 260-350 2	45, 50, 80
Turbots Pleuronectidae spp.	46 \pm 14 18-60 3	86 \pm 28 40-137 3	1.1 \pm 0.4 0.6-1.5 2			1.3 \pm 0.3 1.0-1.5 2	250 \pm 0 250-250 3	3, 15, 111, 119, 127, 138
Whiting Sillaginidae spp.	36 \pm 15 21-50 2	80 \pm 20 60-100 2	6.1 \pm 0.7 5.4-6.9 2					64, 80
Wolffishes Anarhichadidae spp.				350 1		2.0 1	570 1	111
Wrasses Labridae spp.				80 1		3.0 \pm 2.3 0.7-5.3 2		42, 57, 153, 154

¹Average of the means.²Range of the means.³Number of mean values.

LITERATURE CITATIONS continued.

- edulis*) II. Hidratos de carbono. Invest. Pesq. 11(6):33-37.
38. French, R. B., O. D. Abbott, and R. O. Townsend. 1951. Levels of thiamine, riboflavin and niacin in Florida-produced foods. Univ. Florida Bull. 482:1-19.
 39. Fujimaki, M., S. Odagiri, and C. Inagaki. 1953. On the so-called vitamin C of canned bonito in brine. Bull. Jpn. Soc. Sci. Fish. 18:703-708.
 40. Fujita, A., and T. Ebihara. 1939. Über die verteilung des vitamins C in tierischen und pflanzlichen gewebe. II. Biochem. Z. 300:142-147.
 41. Gangal, S. V., and N. G. Magar. 1962. Chemical composition of crabmeat. Indian J. Appl. Chem. 25:133-136.
 42. Giroud, A., C. P. LeBlond, R. Ratismamanga, and E. Gero. 1938. Le taux normal en acide ascorbique. Bull. Soc. Chem. Biol. 20:1079-1087.
 43. Goldbeck, C. G. 1947. Some studies on the content of thiamine and anti-thiamine factor in fishery products. Commer. Fish. Rev. 9(8):13-21.
 44. Gonzalez-Diaz, C., O. Fernandez, and R. O. Cravioto. 1948. Valor nutritivo del pescado fresco que se consume en la Ciudad de Mexico. An. Esc. Nac. Cienc. Biol. Mex. 5:283-290.
 45. Hashimoto, Y., S. Yamada, and T. Mori. 1953. Animal protein factor (APF) and vitamin B₁₂ in marine products—I. Aquatic animals. Bull. Jpn. Soc. Sci. Fish. 19:135-140.
 46. Hastings, W. H., and C. F. Spencer. 1952. Determinations of free and bound ascorbic acid in fishery products. J. Mar. Res. 11:241-244.
 47. Henderson, L. M., H. A. Waisman, and C. A. Elvehjem. 1941. The distribution of pyridoxine (vitamin B₆) in meat and meat products. J. Nutr. 21:589-598.
 48. Higashi, H. 1961. Determination of vitamin D in fish and fish products by chromatography with Japanese acid clay. J. Vitaminol. (Kyoto) 7:215-230.
 49. Higashi, H., S. Hirao, K. Shimizu, J. Yamada, and R. Kikuchi. 1953. Studies on fluctuation of the vitamin A content in fishes—I. Study on Vitamin A in the dogfish, *Squalus suckleyi*. Bull. Jpn. Soc. Sci. Fish. 18:310-318.
 50. Higashi, H., S. Murayama, M. Yanase, and K. Tabei. 1959. The pantothenic acid content of fish and shellfish. Bull. Jpn. Soc. Sci. Fish. 24:770-775.
 51. Higashi, H., S. Hirao, J. Yamada, and R. Kikuchi. 1958. Vitamin contents in the lamprey, *Entosphenus japonicus*. J. Vitaminol. (Kyoto) 4:88-89.
 52. Higashi, H., S. Murayama, and K. Tabei. 1959. Riboflavin contents in the liver of fish. Bull. Jpn. Soc. Sci. Fish. 25:680-686.
 53. Higashi, H., S. Hirao, J. Yamada, and R. Kikuchi. 1962. Studies on the fluctuation of the vitamin A content in fishes—V. Some notes on vitamin A in rainbow trout. Bull. Jpn. Soc. Sci. Fish. 28:715-721.
 54. Higashi, H. 1962. Relationship between processing techniques and the amount of vitamins and minerals in processed fish. In E. Heen and R. Kreuzer (editors), Fish in nutrition, p. 125-131. Fishing News (Books) Ltd., Lond.
 55. Hirao, S., J. Yamada, and R. Kikuchi. 1954. Vitamin A in fish meat—I. Variation in the vitamin A content in fish meat by the anatomical locality. Bull. Jpn. Soc. Sci. Fish. 19:1047-1060.
 56. Hirao, S., S. Murayama, M. Yanase, J. Yamada, R. Kikuchi, and K. Tabei. 1958. Study on the green meat of tuna—I. Quantitative difference of vitamin B group and minerals between the green and the normal meat of precooked tuna. Bull. Jpn. Soc. Sci. Fish. 24:671-675.
 57. Hirao, S., J. Yamada, and R. Kikuchi. 1955. Vitamin A in fish meat—III. Individual fluctuation in the vitamin A content in fish meat. Bull. Jpn. Soc. Sci. Fish. 20:853-861.
 58. Hirao, S., J. Yamada, and R. Kikuchi. 1955. Vitamin A in fish meat—V. Relation between vitamin A and carotenoid pigment in the skin of mackerel. Bull. Jpn. Soc. Sci. Fish. 21:962-966.
 59. Hoar, W. S., and M. Barberie. 1945. Distribution of riboflavin in fresh and processed fish. Can. J. Res. 23E:8-18.
 60. Hoogland, P. L. 1953. The B-vitamins of cod and haddock. Fish. Res. Board Can. Prog. Rep. Atl. Coast Stn. 55:11-14.
 61. Ikeda, S., M. Sato, and R. Kimura. 1963. Biochemical studies on L-ascorbic acid in aquatic animals—II. Distribution in various parts of fish. Bull. Jpn. Soc. Sci. Fish. 29:765-770.
 62. Ikeda, S., and T. Taguchi. 1966. Improved

- assay method and levels of vitamin E in fish tissues. Bull. Jpn. Soc. Sci. Fish. 32:346-351.
63. Ingalls, R. L., J. F. Klocke, J. P. Rafferty, R. E. Greensmith, M. L. Chang, P. I. Tack, and M. A. Ohlson. 1950. Nutritive value of fish from Michigan waters. Mich. Agric. Exp. Stn. Q. Bull. 219, 24 p.
64. Intengan, C. L., L. G. Alejo, I. Concepcion, V. A. Corpus, R. D. Salud, I. Del Rosario, R. Gomez, and J. Henson. 1956. Composition of Philippine foods. V. Philipp. J. Sci. 85:203-213.
65. Jaffe, W. G. 1956. Composition of Venezuelan fishes. Arch. Venez. Nutr. 7:163-166.
66. Joshi, S., F. Master, and N. G. Magar. 1953. Nutritive value of some Bombay fish. Part I. Distribution of non-protein nitrogen extractives, and thiamin, riboflavin and niacin. Indian J. Med. Res. 41:431-439.
67. Kühnau, J. 1956. Tabellen der chemischen Zusammensetzung von fischen. I. Vitamine. Arch. Fischereiwiss. 7(suppl.), 28 p.
68. Kelley, C. E., and A. W. Harmon. 1972. Method of determining carotenoid contents of Alaska pink shrimp and representative values for several shrimp products. Fish. Bull., U.S. 70:111-113.
69. Khan, A. H., and S. A. Haq. 1958. Studies on marine edible (Teleostii) fishes. Part I.—Distribution of oil and vitamin A in the skin, flesh and liver of edible fishes of Karachi waters. Pak. J. Sci. Ind. Res. 1:309-311.
70. Khorana, M. L., M. L. Sarma, and K. V. Giri. 1942. Investigations on the food value of fish and other marine products. Part I. The antipellagra vitamin (nicotinic acid). Indian J. Med. Res. 30:315-318.
71. Klocke, J. F., T. Porter, P. I. Tack, E. Leffler, N. S. Henry, and R. Nitchals. 1946. Nutritive value of fish from Michigan waters. I. Nicotinic acid of lake herring, carp, common suckers, and burbot. Food Res. 11:179-186.
72. Klocke, J. F., P. I. Tack, M. A. Ohlson, R. Nitchals, E. Leffler, and N. S. Henry. 1947. Nutritive value of fish from Michigan waters. II. Thiamin of lake herring, carp, common sucker, burbot, and sinelt. Food Res. 12:36-43.
73. Kodicek, E. 1940. Estimation of nicotinic acid in animal tissues, blood and certain foodstuffs. 2. Applications. Biochem. J. 34:724-735.
74. Komata, Y., Y. Hashimoto, and T. Mori. 1956. B-vitamins in marine products and their changes in the processing and storage—I. Canned mackerel and tuna in brine. Bull. Jpn. Soc. Sci. Fish. 21:1236-1240.
75. Komata, Y., Y. Hashimoto, and T. Mori. 1957. B-vitamins in marine products and their changes during the processing and storage—II. Canned baby clam. Bull. Jpn. Soc. Sci. Fish. 23:44-46.
76. Kon, S. K. 1949. Preformed vitamin A in northern krill. Proc. Biochem. Soc., 28th meeting, p. 31-32.
77. Lane, R. L., E. Johnson, and R. R. Williams. 1942. Studies of the average American diet. I. Thiamine content. J. Nutr. 23:613-624.
78. Le Cornu, L. R. 1949. High food value of fish. Fish. Newsl. 8(6):10-11.
79. Leong, P. C. 1940. Nicotinic acid content of foods. J. Malaya Branch, Br. Med. Soc. 4(3):261-278.
80. Leung, W. W. 1972. Food composition table for use in east Asia. Natl. Inst. Arthritis, Metabolism Digestive Diseases, N.I.H., 334 p.
81. Leung, W.-T. W., R. K. Pecot, and B. K. Watt. 1952. Composition of foods used in Far Eastern countries. U.S. Dep. Agric., Agric. Handb. 34, p. 35-38.
82. Leung, W. W. 1968. Food composition table for use in Africa. U.S. Dep. Health Educ. Welfare, 306 p.
83. Leung, W. W. 1961. Food composition table for use in Latin America. A research project sponsored jointly by: Inst. Nutr. Central America and Panama, Guatemala City, C. A. and The Interdepartmental Commission on Nutrition for National Defense, N.I.H.
84. Lieck, H., and H. Sondergaard. 1958. The content of vitamin B₆ in Danish foods. Int. Z. Vitaminforsch. 29:68-77.
85. Lontoc, A. V., and O. N. Gonzalez. 1968. Vitamin B₁₂ content of some Philippine foods. Philipp. J. Nutr. 21(3):163-171.
86. Loughlin, M. E., and A. E. Teeri. 1960. Nutritive value of fish. II. Biotin, folic acid, pantothenic acid, and free amino acids of various salt-water species. Food Res. 25:479-483.
87. Ludány, G. V. 1936. C-vitamin in Subwasserfischen und Subwasserkrebsen. Biochem. Z. 284:108-110.
88. Lunde, G., V. Aschehoug, and H. Kringstad. 1937. Fish as a source of vitamins A and D. Herring, brisling, and their products. Ind. Eng. Chem. 29:1171-1175.
89. Lunde, G. 1939. Neuere forschungen über die vitamine in fischen und fischprodukten. Angew. Chem. 52(32):521-532.
90. Lunde, G., H. Kringstad, and A. Olson. 1938. Untersuchungen über den gehalt an vitamin B₁ und B₂. I. Mat.-Naturk. Klass. 7:3-51.
91. Marks, A. L., and H. W. Nilson. 1946. Thiamin and riboflavin in baked and simmered oysters. Commer. Fish. Rev. 8(12):12-13.
92. Martinek, W. A., and C. G. Goldbeck. 1947. Nutritive value of baked croaker. Commer. Fish. Rev. 9(4):9-13.
93. Martinez-Tamayo, J. A. 1964. Estudio de la vitamina B₆ y su determinacion espectrofotometrica en alimentos. An. Brom. 16(3):239-262.
94. Maxwell, B. E. 1952. The distribution of vitamin B₁₂-active substances in some marine invertebrates of British Columbia. J. Fish. Res. Board Can. 9:164-168.
95. McCance, R. A., and E. M. Widdowson. The composition of foods. Med. Res. Council, Spec. Rep. Ser. 297, 252 p.
96. McCance, R. A. 1944. The chemistry of growth and the food value of the common eel (*Anguilla anguilla* (L.)). Biochem. J. 38:474-480.
97. McVicar, R. W., and G. H. Berryman. 1942. Nicotinic acid in foods. J. Nutr. 24:235-243.
98. Miller, C. D. 1957. Nutritive values of some Hawaiian foods in household units and common measures. Hawaii Agric. Exp. Stn. N. 52, p. 3-20.
99. Miwa, K., Y. Sato, and E. Kinoshita. 1958. Studies on the utilization of migratory fish.—II On the constituent parts of body of saury and their chemical component. Bull. Hokkaido Reg. Fish. Res. Lab. 19:66-71.
100. Miyake, M., and K. Hiyashi. 1961. Vitamin B group in the extracts of Mollusca—I. On vitamin B₆. Bull. Jpn. Soc. Sci. Fish. 27:458-460.
101. Mori, T., Y. Hashimoto, and Y. Maeda. 1954. Animal protein factor (APF) and vitamin B₁₂ in marine products—IV. Variations in the vitamin B₁₂ content of marine animals in the spoilage. Bull. Jpn. Soc. Sci. Fish. 19:991-996.
102. Mori, T., Y. Hashimoto, and Y. Komata. 1956. B-vitamins content in the muscle of fish. Bull. Jpn. Soc. Sci. Fish. 21:1233-1235.
103. Mori, T., S. Konosu, and S. Miyagawa. 1957. Choline content of the muscle of aquatic animals. Bull. Jpn. Soc. Sci. Fish. 23:282-285.
104. Munsell, H. E. 1942. Riboflavin content of some common foods. Food Res. 7:85-95.
105. Murayama, S., and K. Tabei. 1956. Studies on the vitamin B group (B₁, B₂, B₆ and B₁₂) in the pyloric appendages of fishes. Bull. Jpn. Soc. Sci. Fish. 22:136-141.
106. Murayama, S., M. Yanase, and K. Tabei. 1959. Vitamin B contents of tissues of red salmon (*Oncorhynchus nerka*). Fish. Res. Bull. 25:67-69.
107. Nazir, D. J., and N. G. Magar. 1963. Chemical composition of mackerel (*Rastrelliger canagurta*) and changes in the nutritive value during storage. Indian J. Technol. 1:247-249.
108. Novak, A. F., E. A. Fieger, and M. E. Bailey. 1956. Vitamin content of fresh, processed shrimp. Food Freezing 18:9-10.
109. Nguyen-Thi-Lau, and C. Richard. 1959. Le poisson dans l'alimentation du Vietnam. Rev. d'Elev. Med. Vet. 12:313-324.
110. Obata, Y., and T. Yamanishi. 1952. Chemical studies on the substance of fish smell V. Aroma of cooked fish. Jpn. Soc. Sci. Fish. Bull. 17:326-328.
111. Orr, M. L. 1969. Pantothenic acid, vitamin B₆ and vitamin B₁₂ in foods. U.S. Dep. Agric., Home Econ. Res. Rep. 36:13-26.
112. Palad, J. G., I. C. Abdon, A. V. Lontoc, L. B. Dimaunahan, E. C. Eusebio, and N. Santiago. 1964. Nutritive value of some foodstuffs processed in the Philippines. Philipp. J. Sci. 93:355-384.
113. Pandit, A. R., and N. G. Magar. 1972. Chemical composition of Sepia orientalis and Loligo vulgaris. Fish. Technol. 9:122-125.
114. Peden, J. D. 1973. Heavy metals in Somerset marine organisms. Mar. Pollut. Bull. 4(1):7-9.
115. Polansky, M. M., and E. W. Toepfer.

1969. Vitamin B₆ components in some meats, fish, dairy products, and commercial infant formulas. *J. Agric. Food Chem.* 17:1394-1397.
116. Pradhan, V., and R. G. Chitre. 1951. Thiamine and riboflavin content in the tissues of some salt water fishes in Bombay. *Indian J. Med. Sci.* 5:645-648.
117. Proctor, B. E., S. A. Miller, S. A. Goldblith, E. L. Wick, E. R. Pariser, G. M. Sapers, and M. Solberg. 1961. The nutritive value of Maine sardines. I. Chemical composition. *J. Food Sci.* 26:283-287.
118. Prudhomme, M. 1960. Les escargots. *Rev. Med. Vet.* 23:456-461.
119. Pyke, M. 1939. The distribution of vitamin B₁ in foods as determined by chemical analysis. *J. Soc. Chem. Ind.* 58:338-340.
120. Pyke, M., and M. D. Wright. 1941. Vitamin content of salmon. *Nature (Lond.)* 147:267.
121. Robbins, W. J., A. Hervey, and M. E. Stebbins. 1951. Further observations on Euglena and B₁₂. *Bull. Torrey Botanical Club* 78:363-375.
122. Rudra, M. N. 1936. Studies in vitamin C. II. The vitamin C contents of the liver and muscle of some Indian fresh-water fish. *J. Indian Chem. Soc.* 13:740-742.
123. Sagara, J., and M. Yanase. 1958. Vitamin B₁₂ as a factor in the environment of bivalves—I. Seasonal and environmental variation of vitamin B₁₂ content in *Tapes japonica* and *Meretrix meretrix lusoria*. [In Jpn., Engl. summ.] *Bull. Jpn. Soc. Sci. Fish.* 24:456-461.
124. Saha, K. C. 1941. Nicotinic acid content of fish. *Ann. Biochem. Exp. Med. (Calcutta)* 1:75-78.
125. Sautier, P. M. 1946. Thiamine assays of fishery products. *Commer. Fish. Rev.* 8(2):17-19.
126. Sawant, P. L., and N. G. Magar. 1961. Nutritive value of canned fish: Effect of canning, storage & antioxidants. *J. Sci. Ind. Res.* 20D:313-316.
127. Schroeder, H. A. 1971. Losses of vitamins and trace minerals resulting from processing and preservation of foods. *Am. J. Clin. Nutr.* 24:562-573.
128. Shaikhmahmud, F., and N. G. Magar. 1957. Studies in the nutritive value of Bombay prawns: Part I—Chemical composition of prawns. *J. Sci. Ind. Res.* 16A:44-46.
129. Shaikhmahmud, F. S., and N. G. Magar. 1961. Studies in the nutritive value of Bombay prawns: Part II—Chemical composition & nutritional constituents of Penaeid prawns. *J. Sci. Ind. Res.* 20D:157-158.
130. Shimada, K., and E. Tanaka. 1936. Distribution of inositol in fish. *Bull. Jpn. Soc. Sci. Fish.* 4:305-306.
131. Shimidu, Y., M. Yano, and S. Higasa. 1959. Biochemical studies on the growth and maturation of fish. X. Some notes on riboflavins in the skin of tunas and marlins. *Rep. Nankai Reg. Fish. Res. Lab.* 11:24-31.
132. Sidwell, V. D., P. R. Foncannon, N. S. Moore, and J. C. Bonnet. 1974. Composition of the edible portion of raw (fresh or frozen) crustaceans, finfish, and mollusks. I. Protein, fat, moisture, ash, carbohydrate, energy value, and cholesterol. *Mar. Fish. Rev.* 36(3):21-35.
133. Sidwell, V. D., D. H. Buzzell, P. R. Foncannon, and A. L. Smith. 1977. Composition of the edible portion of raw (fresh or frozen) crustaceans, finfish, and mollusks. II. Macroelements: Sodium, potassium, chlorine, calcium, phosphorus, and magnesium. *Mar. Fish. Rev.* 39(1):1-11.
134. Sidwell, V. D., A. L. Loomis, K. J. Loomis, P. R. Foncannon, and D. H. Buzzell. 1978. Composition of the edible portion of raw (fresh or frozen) crustaceans, finfish, and mollusks. III. Microelements. *Mar. Fish. Rev.* 40(9):1-19.
135. Sinnhuber, R. O., T. C. Yu, T. C. Yu, and N. L. Karrick. 1956. Variation in proximate composition of right and left fillets of rockfish (*Sebastes pinniger*) and Dover sole (*Microstomus pacificus*) *Commer. Fish. Rev.* 18(12):24-27.
136. Sondergaard, H. 1959. Statens vitaminlaboratorium undersogelser af B₆-vitaminindholdet i danske naeringsmidler. *Stat. Husholdn. Fag. Med.* 1-2:1-7.
137. Sreenivasamurthy, V., M. Swaminathan, and V. Subrahmanyam. 1954. Vitamin B₁₂ content of some animal foods. I. *Bull. Central Food Tech. Res. Inst. Mysore* 9:33-38.
138. Swaminathan, M. 1946. Vitamin B₁ (thiamin) content of Indian foodstuffs. *Indian J. Med. Res.* 34:289-297.
139. Swaminathan, M. 1944. Nicotinic acid content of Indian foodstuffs. *Indian J. Med. Res.* 32:39-46.
140. Taarland, T., E. Mathiesen, Ø. Øusthus, and D. R. Brackkan. 1958. Nutritional values and vitamins of Norwegian fish and fishproducts. [In Norw., Engl. summ.] *Tidssk. Hermetikind.* 44:405-412.
141. Tack, P. I., R. Ingalls, J. F. Klocke, H. Baeder, D. C. Cederquist, E. Musser, and M. A. Ohlson. 1947. Preparation, utilization, and vitamin concentrations of the burbot, *Lota lota maculosa* (Lesueur). *Mich. Q. Bull.* 29:286-293.
142. Takada, K., and J. Nishimoto. 1958. Studies on the choline in fish—I. Content of choline and the similar substances in fishes. *Bull. Jpn. Soc. Sci. Fish.* 24:632-635.
143. Tarr, H. L. A., B. A. Southcott, and P. W. Ney. 1950. Vitamin B₁₂—Active substances in fish products. *Food Technol.* 4:354-357.
144. Teeri, A. E., M. E. Loughlin, and D. Josselyn. 1957. Nutritive value of fish. I. Nicotinic acid, riboflavin, vitamin B₁₂ and amino acids of various salt-water species. *Food Res.* 22:145-150.
145. Teply, L. J., F. M. Strong, and C. A. Elvehjem. 1942. The distribution of nicotinic acid in foods. *J. Nutr.* 23:417-423.
146. Thomas, M. H., and D. H. Calloway. 1961. Nutritional value of dehydrated foods. *J. Am. Diet. Assoc.* 39:105-116.
147. Waisman, H. A., L. M. Henderson, J. M. McIntire, and C. A. Elvehjem. 1942. The effect of enzymatic digestion on the pantothenic acid content of meats determined by the microbiological method. *J. Nutr.* 23:239-248.
148. Wentworth, J., and H. Lewis. 1958. Riboflavin, niacin, and thiamine in Apalachicola Bay oysters. *Food Res.* 23:194-197.
149. Wertz, A. W., B. W. McKay, K. O. Eslesen, and J. O. Holmes. 1944. The effect of institutional cooking methods on the vitamin content of food. II. Thiamine, riboflavin, nicotinic acid and pantothenic acid content of fish. *Mass. Agric. Ext. Stn. Prog. Notes* 3, p. 1-4.
150. Willimott, S. G. 1949. Malayan food composition table. *Dep. Agric. Fed. Malaya Sci. Ser. No.* 23, 24 p.
151. Yamada, J. 1962. Vitamin A in *Entosphenus tridentatus* Gairder. [In Jpn., Engl. summ.] *Bull. Tokai Reg. Fish. Res. Lab.* 33:33-34.
152. Yamadori, M. 1949. Determination of standard vitamin contents of Japanese food stuffs. *Kitasato Arch. Exp. Med.* 22:41-46.
153. Yanase, M. 1958. The vitamin B₆ contents of fish liver. [In Jpn., Engl. summ.] *Bull. Tokai Reg. Fish. Res. Lab.* 22:53-56.
154. Yanase, M. 1953. Studies on the vitamin B₁₂ content of shell fish. *Bull. Jpn. Soc. Sci. Fish.* 18:636-638.
155. Yanase, M. 1963. Pantothenic acid and vitamin B₁₂ contents assessed for the rainbow trout during maturation. *Bull. Jpn. Soc. Sci. Fish.* 29:1024-1026.
156. Yanase, M. 1953. Studies on the vitamin B₁₂ of aquatic animals—II. The vitamin B₁₂ content of fish (2). *Bull. Jpn. Soc. Sci. Fish.* 18:629-635.
157. Yanase, M. 1956. The vitamin B₆ content of fish meat. *Bull. Jpn. Soc. Sci. Fish.* 22:51-55.

MFR Paper 1355. From *Marine Fisheries Review*, Vol. 40, No. 12, December 1978. Copies of this paper, in limited numbers, are available from D822, User Services Branch, Environmental Science Information Center, NOAA, Rockville, MD 20852. Copies of *Marine Fisheries Review* are available from the Superintendent of Documents, U.S. Government Printing Office, Washington, DC 20402 for \$1.10 each.

Review of Oxygen Depletion and Associated Mass Mortalities of Shellfish in the Middle Atlantic Bight in 1976

FRANK W. STEIMLE and CARL J. SINDERMANN

ABSTRACT—In summer and autumn of 1976, mass mortalities of shellfish occurred in a 165-km long corridor of severe oxygen depletion paralleling the New Jersey coast from 5 to 85 km from shore. Mortalities of surf clams, *Spisula solidissima*, the most severely affected species, were estimated in excess of 140,000 t. Alteration of normal migration patterns of lobsters and several species of finfish was also noted. A series of anomalous meteorological and hydrological events (particularly early warming of surface waters resulting in early thermocline development, and a massive shelf-wide phytoplankton bloom) superimposed on an already stressed coastal area, was considered to be responsible. The occurrence is particularly significant because the continental shelf of the Middle Atlantic Bight, from Cape Cod to Cape Hatteras on the east coast of the United States, contains the largest known stocks of ocean shellfish of any comparable coastal area of North America.

INTRODUCTION

Mass mortalities in the sea are relatively common events and always attract the interest of the scientific community as well as the public. This interest may be based on a concern for the loss of a fishery resource, or the nuisance or health problems created by the decaying animals. Very often it is difficult to identify the cause of mass mortalities because most investigations begin after the fact, so the conditions which lead to the mortalities may have been altered or dissipated by the time studies begin. The majority of mass mortalities are very localized, often confined to a particular bay or estuary, but a few can be widespread, sometimes affecting hundreds of square kilometers of ocean.

The recognized causes of the mass mortalities have been physical (vulcanism, rapid or extreme temperature changes, storms, seaquakes, and

strandings), chemical (extreme salinity or pH changes, oxygen depletion, toxic chemicals, and hydrogen sulfide formation), biological (disease and toxic or massive algal blooms), or combinations of the above. Many mass mortalities have been reported in the scientific literature since the 19th century and fossil evidence of mass mortalities exists. Brongersma-Sanders (1957) has compiled the most comprehensive review to date; other reviews include those of Sinderman (1970, 1976).

An environmental event of heroic proportions, leading to mass mortalities of many marine species in a 12,000 km² area of the continental shelf off the Middle Atlantic coast of the United States occurred during July through October 1976. Investigators were able to detect conditions that were lethal to marine life; these conditions were extreme oxygen depletion and hydrogen sulfide formation in bottom waters. First reports of this developing en-

vironmental problem reached the scientific community during the weekend of 4 July. Sport divers, lobstermen, and trawler fishermen had observed and reported dead and dying marine organisms, both fish and invertebrates, on fishing reefs and wrecks, and on fishing grounds, off the north-central New Jersey coast, south of New York City. Within a few weeks the mortalities were reported in areas extending southward some 85 km and well out on the continental shelf.

A series of survey cruises was initiated by the National Marine Fisheries Service's Northeast Fisheries Center, Sandy Hook, N.J., to examine the extent of the problem, to assess the damage, and investigate possible causes. Oxygen deficient bottom water, sometimes with zero dissolved oxygen levels, was found for a north-south distance of 165 km in a zone or corridor from 5 to 85 km off the New Jersey coast.

In the central part of this zone, oxygen values were zero, and hydrogen sulfide was detected below the thermocline. Oxygen depletion persisted until October, when lower surface temperatures and mixing, after disappearance of the thermocline, gradually reoxygenated the bottom water.

Mortalities of fish, lobsters, molluscan shellfish, and other benthic invertebrates were observed. The sedentary forms, surf clams, ocean quahogs, and sea scallops, suffered the greatest mortalities. From almost continuous surveys, it was estimated that 69 percent of the surf clam population off the New Jersey coast, representing some 143,000 t of meats, had been destroyed by October, with significant but lesser mortalities of ocean quahogs and sea scallops. Lobster catches were reduced by 30 percent during the period. The New Jersey coast was declared a resource disaster area in November by the

Frank W. Steimle and Carl J. Sindermann are with the Sandy Hook Laboratory, Northeast Fisheries Center, National Marine Fisheries Service, NOAA, Highlands, NJ 07732.

Federal government because of this event.

The problem was complicated by the public's premature conclusion that ocean disposal of pollutants, particularly sewage sludge dumping (over 5 million tons were dumped in 1976), 20 km from the coast, was responsible for this oxygen depletion situation and associated mortalities. Concern was also expressed that this catastrophic event might be repeated, possibly even annually.

The extent and duration of the problem and tremendous impacts on the fisheries and businesses in the area stimulated a massive investigation effort by State, Federal, and private groups or agencies. An interagency group was created to coordinate research efforts and finally to draw all available data together by holding a series of workshops, during November 1976. The results of these workshops, which are for the most part preliminary analyses of available data were published as an interim report (National Marine Fisheries Service, 1977).

From these workshops a large amount of data has been assembled and a causal hypothesis developed, centering on a combination of unique atmospheric, hydrographic, and biological events, which occurred in a coastal area already stressed by human organic loading.

The major environmental disturbances of 1976 in the Middle Atlantic Bight may prove to be one of the best-documented examples that we have of a mass mortality in the sea and its short- and long-term impacts on resource and foodchain species. Scientific studies are continuing, especially since the possibility of repetition of the event, at some level of intensity, exists for future years.

This paper is a review of the situation based partially on the workshop reports and includes subsequent data and assessments.

EXTENT OF OXYGEN DEPLETION

One major aspect of the 1976 investigation was the monitoring of the levels and distribution of dissolved




Figure 1.—The bottom oxygen levels in New York Bight at period of greatest distribution, late September 1976.

oxygen in the sub-thermocline bottom waters. Sampling in mid-July found depressed dissolved oxygen (D.O.) values, some below the level of detection by standard Winkler procedure, in an area 3-35 km off Barnegat Inlet, 115 km south of New York City (Fig. 1). Values of less than 2 ppm were found between Long Branch to north of Atlantic City, a coastal distance of 105 km. Trawl surveys collected dead epibenthic invertebrates and stressed surf clams in this zone and noted an absence of the normal finfish population known to inhabit the area in the summer.

By early August, prior to Hurricane Belle, which passed the New Jersey coast on 10 August, the anoxic area had moved or expanded southward, with the center of the oxygen-depleted area being found between Barnegat Inlet and Atlantic City, a distance of 70 km on the central New Jersey coast. Extremely high levels of hydrogen sulfide (to 1.76 mg/l) were also detected near the center of the depleted area. The hydrogen sulfide was present up to 15 m from the bottom (35 m deep), but not above the thermocline (Draxler and Byrne, 1977).

Hydrogen sulfide was also evident in an apparent upwelling of anoxic bottom water along very restricted portions of the immediate shoreline in central New Jersey. Hundreds of fish of several species, including sharks, were trapped along the beach and killed. A period of strong westerly winds, pushing the in-shore surface waters offshore, was thought responsible.

The hurricane, from which many had hoped for relief, did not significantly alter the situation. Immediately after its passage, resurveys of stations off Atlantic City which had been surveyed just prior to the storm found some possible coastal mixing or an offshore shift, resulting in the less than 2 ppm D.O. area moving from 3 km off the coast to 25 km offshore as the only apparent effect. This was only temporary, because a second resurvey, 5 days later, indicated the anoxic water mass had resumed most of its prehurricane distribution, with further movement or expansion south-southeast (Steimle, 1977a).

By mid-September, the anoxic area (defined here as the area where bottom water D.O. values were less than 2 ppm), reached its greatest known distribution, covering approximately half the New York Bight, including a tongue off Long Island, and extending southward to the Maryland State border (Fig. 1).

By the first week in October, surveys found that the thermocline was apparently decaying because bottom D.O. concentrations inshore, out from the coast to 40 km, were increasing to nonhazardous levels. By early November, no trace of oxygen depletion was evident and the 1976 oxygen depletion phenomenon had evidently ended.

RELATED METEOROLOGICAL AND HYDROLOGICAL EVENTS

Data from intensive field operations in the Middle Atlantic Bight during the critical period, combined with meteorological observations, were examined in the interagency workshops. It was the general conclusion of the workshops (National Marine Fisheries Service, 1977) that large-scale meteorological and oceanographic phenomena were

involved in production of the anoxic zone which resulted in mortalities. The hypothesis which was developed focused on a somewhat unique combination of anomalous environmental events superimposed on a marginal coastal area, which has been made eutrophic by man's input of organic material.

Meteorological events included: 1) High February-March temperatures with peak river runoff in February instead of April; 2) reduction of cyclonic storm activity during the summer to less than half the 25-year average; and 3) a period of 4-6 weeks in June-July with persistent south or southwest winds (Diaz, 1976).

Physical oceanographic events included: 1) Early (February-March) warming of surface waters and development of the thermocline (thermal stratification usually begins in April and reaches a maximum in August); and 2) early onset of decline in subsurface dissolved oxygen values (January rather than March). Bottom water D.O. values in May 1976 off New Jersey were as low as they usually are in July (Armstrong, 1977).

Biological oceanographic events included: 1) A massive bloom of the dinoflagellate *Ceratium tripos* over much of the Middle Atlantic Bight, but particularly concentrated in the New York Bight. The bloom began in February, persisted at least until July, and was concentrated at and just below the thermocline (Malone, 1977).

Since the Middle Atlantic Bight has been the focus of intensive research by elements of the National Oceanic and Atmospheric Administration for several years, a number of earlier investigations had been conducted. This historical data base enabled comparison with conditions which existed in previous years. Based on such comparisons, an explanation of the events of 1976 could be summarized in several steps: 1) Oxygen demand from a declining phytoplankton bloom was superimposed on a shallow shelf area (New Jersey coast) already characterized by reduced dissolved oxygen in an average summer; 2) this organically rich oxygen demanding water was sealed off early

in spring by the early onset of a thermocline; 3) water mass movement was reduced to a minimal southward flow of bottom water; and 4) cyclonic storm activity during the entire period was abnormally low. With these factors, the ingredients of disaster to marine animals were present.

FISH AND SHELLFISH STOCKS OF THE MIDDLE ATLANTIC BIGHT

Fish and shellfish populations of the Middle Atlantic Bight are abundant and important to the nation's economy. Oceanic species of bivalve mollusks, particularly surf clams, ocean quahogs, (*Arctica islandica*), and scallops (*Placopecten magellanicus*) are more numerous here than in any comparable coastal area in the United States. Surf clams harvested from the Middle Atlantic Bight constitute over 50 percent of total landed weight of molluscan shellfish in the United States; the fishery for ocean quahogs is expanding rapidly, and populations of sea scallops are fished regularly in deeper waters of the Bight.

The National Marine Fisheries Service has conducted surveys of surf clam, ocean quahog, and scallop distribution and abundance in the Middle Atlantic Bight for a number of years. The most recent surveys for surf clams, ocean quahogs, and sea scallops were in April 1976. Total estimated biomass of offshore surf clams in the Bight was 875,000 t of meats, with the New Jersey sector containing 207,000 t. Total estimated biomass of ocean quahogs in the Bight was 2,450,000 t of meats, with the New Jersey sector containing 818,000 t (Ropes and Chang, 1977). Biomass estimates for sea scallops in the Bight are not available, but much of the stocks are composed at present of a single strong year class (1972). Scallops occupy about 11,500 km² of the shelf off New Jersey (MacKenzie, 1977).

Finfish species presently of significance in the Middle Atlantic Bight include scup (*Stenotomus chrysops*), summer flounder (*Paralichthys dentatus*), bluefish (*Pomatomus saltatrix*),

striped bass (*Morone saxatilis*), mackerel (*Scomber scombrus*), black sea bass (*Centropristis striata*), and weakfish (*Cynoscion regalis*). A number of these species are taken by recreational as well as commercial fishermen, often with the recreational catch predominating. Some of the species exist as year-round local populations, while others, such as summer flounder and bluefish, migrate vertically to and from the coast, or laterally north and south through the Bight. A few species (mackerel and silver hake, *Merluccius bilinearis*) have until recently been exploited heavily by foreign distant-water fleets. Most of the Middle Atlantic finfish stocks of interest to U.S. fishermen (other than mackerel) have not declined drastically in recent decades, and increased landings characterize species such as bluefish, striped bass, and weakfish since 1970 (McHugh, 1977).

The National Marine Fisheries Service (and its predecessor agencies) has also conducted trawling surveys for demersal fish in the Middle Atlantic Bight for the past decade to provide a basis for continuing stock assessment. Additionally, recreational fish surveys have been carried out during the past 3 years. Landing figures for important recreational and commercial species are summarized in Figure 2.

EFFECTS ON SHELLFISH AND OTHER BENTHIC POPULATIONS

Beginning in late July 1976, assessment of the impact of the anoxic event on the surf clam stocks began. Signs of stressed surf clams were noted by divers as early as the weekend of 4 July. These were clams that were not embedded in the sediment but were lying free on the surface. Several later trawl surveys also found live, but gaping clams. The first specific surf clam dredging survey was completed by the end of July. Mortalities ranging between 0 and 56 percent were found in a restricted area off Barnegat Inlet. A second survey, in early August, found an average mortality of 10 percent in clam stocks in




Figure 2.—Landings of recreational and commercial species in the Middle Atlantic Bight in 1970 (National Marine Fisheries Service, 1973).

the impacted area. The normal mortality is 2 percent (Ropes and Chang, 1977).

Subsequent expanded resurveys in September and October found that the average mortality had risen to 100 percent at some stations in a 12,000 km² sector off New Jersey. It was estimated that this represented, by October 1976, a loss of 143,000 t of surf clam meats, or about 69 percent of the offshore surf clam stocks of New Jersey, and 16 percent of the estimated total Middle Atlantic Bight population of the species (Fig. 3). Of the coastal surf clam stocks (those within 5 km of shore) an estimated 1,700 t were killed in an area south of Beach Haven—representing about 5 percent of the total inshore coastal population (Ropes and Chang, 1977). Because July is the normal spawning season for surf clams, the impact on future stocks may also be severe.

Mortalities were observed also in New Jersey's ocean quahog population, an increasingly valuable resource species, which is usually found in deeper water than the surf clam. In early August, mortalities for this species were less than 1 percent. Mortalities increased in September to almost 8 percent, with a high of 40 percent at some individual stations. The loss to New Jersey stocks of ocean quahogs was about 6,600 t, or less than 1 percent of the stocks in that sector of the coast (Fig. 4) (Ropes and Chang, 1977).

Sea scallops, which also occur in deeper water than surf clams, were affected by the anoxic event. Scallops occupy an area of 11,500 km² off the New Jersey coast; of this, 4,300 km² were within the anoxic zone, and an estimated 10 percent of the population was killed in a zone 35-55 m deep (Fig. 5) (MacKenzie, 1977).

The lobster, *Homarus americanus*, industry off New Jersey suffered. Some of the inshore stocks were killed, and the annual shoreward migration of offshore stocks was interrupted. During the months of June through September, normally the most productive months of the year, landings declined an average 30 percent compared with the same




Figure 3.—Distribution of surf clam mortalities in New York Bight.

period in 1975. The inshore pot fishery, which operates within 20 km of shore, was most severely affected. Lobstermen stated that few offshore migrants entered the fishery in 1976 (Halgren, 1977).

Other benthic populations were affected by the anoxic water. Effects on the benthic infauna were most noticeable in the H₂S zone, with reduction in numbers of species and numbers of individuals. Species to species variability in survival was noted, with a number of polychaetes and sea anemones quite resistant to prevailing extreme environmental conditions. Among the benthic megafaunal species affected were rock

crabs (*Cancer irroratus*, *C. borealis*), mud shrimp (*Axiu serratus*), mantis shrimp (*Platysquilla enodus*), starfish (*Asteria* sp.), moon snail (*Lunatia heros*), sea cucumbers, (*Thyone* sp.), and sand dollars (*Echinarachnius parma*). Many of the more mobile crustaceans were able to avoid the developing anoxia, but not all of them (Radosh et al., 1977; Steimle, 1977b).

Oxygen depletion in bottom waters may prove to be a common feature in localized areas where ocean shellfish occur. Drastic changes in abundance of species such as the surf clam, which has been observed previously, may in some instances be caused by anoxia. It may

be too that the presence of anoxic bottom waters may influence post-larval settling and survival of clams, even eliminating set of a particular year class in certain areas.

The survival of marine invertebrates, including shellfish, in oxygen-depleted waters, and in the presence of hydrogen sulfide, has been examined, and additional experiments are being conducted by the National Marine Fisheries Service. Earlier studies (Theede et al., 1969; Davis, 1975) and preliminary results of current studies indicate survival of clams for periods of several weeks in water which approaches zero D.O., but shorter survival in hydrogen sulfide environments. These experimental findings agree with field observations in 1976. Surf clams began showing signs of stress early in July, but mortalities were not reported until the middle of July. The extended period of anoxia, combined with hydrogen sulfide, resulted in 100 percent mortality by October at stations in the most severely affected zone along the central New Jersey coast, but lesser mortality in peripheral areas of less severe oxygen depletion. The majority of the mortalities occurred in July and August, as evidenced by mostly clapper (meatless paired shells) bivalves collected after September.

EFFECTS ON FISH POPULATIONS

Even though some mortalities of finfish were reported early in the event, it seems that the principal effect of oxygen depleted waters below the thermocline was to modify the normal movements and migrations of a number of species. Summer flounder were crowded within a narrow coastal zone or in estuaries, where they were readily available in large numbers to fishermen. Bluefish, which normally migrate northward through the area in summer, were shown by results of tagging studies to have reversed their direction, and moved southward (Freeman and Turner, 1977a, 1977b; Festa, 1977).

Demersal species apparently abandoned the oxygen-minimum and hydrogen sulfide area completely. Several trawling surveys through the area in August, September, and October dis-




Figure 4.—Distribution of ocean quahog mortalities in New York Bight.

closed almost complete absence of any demersal fish species—an unusual event when compared with similar trawling surveys conducted in the Middle Atlantic Bight since 1968. Very few dead fish were brought up in the nets during the 1976 surveys, although dead invertebrates were common (Azarovitz et al., 1977).

There is genuine concern, however, that survival of the 1976 year class of demersal spawning fish may have been severely affected in the anoxic zone (Smith, pers. commun.¹).

¹Smith, W. National Marine Fisheries Service, NOAA, Highlands, N.J. Pers. commun.

DISCUSSION

There are, of course, other coastal areas in the world where extreme oxygen depletion in bottom waters is a frequent, sometimes even an annual, event.

Marine fish kills related to oxygen depletion and hydrogen sulfide buildup have been reported in warm shallow estuaries (May, 1973) and in areas of upwelling and mass production of plankton, e.g., off South America and Africa (Brongersma-Sanders, 1957; Theede et al., 1969).

A coastal upwelling region famous for its low oxygen, hydrogen sulfide production, and periodic mortalities, is




Figure 5.—Distribution of scallop mortalities in New York Bight.

off the southwest coast of Africa, in and near Walvis Bay. Scientific records of mortalities, summarized by Brongersma-Sanders (1947, 1957), extend back to 1837. Dead and dying fish, cephalopods, and bivalves have been observed with great frequency in December and January in the sea and on the beaches between lat. 21°S and 25°S. The sea bottom of the region is highly organic, with high H_2S content and anoxic bottom waters. Mass mortalities of fish are more severe in some years than in others, and are often preceded by red to brown discoloration of the sea from algal blooms. The anoxic area involved is approximately 14,000

km², but, interestingly, there is a narrow coastal strip about 5 km wide, extending to a depth of 40 m, where sea life is normal and hydrogen sulfide does not occur (Copenhagen, 1953). Similar mass mortalities of marine animals in a zone of upwelling have been reported by Falke (1950) from Concepcion Bay, Chile.

Mass mortalities, particularly of benthic fauna, have occurred in the deeper basins of the Baltic, where anaerobic conditions may persist for as long as 4 years (Segerstråle, 1969). Total absence of oxygen, beginning in 1957, caused the deeper parts of the Gotland, Gdansk, and Bornholm Ba-

sins to become lifeless deserts in 1958-59. The total area affected was estimated at 33,000 km². The stagnation was broken in 1962 by a strong inflow of saline water from the Kattegat. It is significant that great amounts of nutrients accumulated during the stagnation period; these were brought to the surface in 1962, resulting in an enormous increase in plankton populations. A similar event had occurred in the early 1930's (Kalle, 1943; Meyer and Kalle, 1950). The most recent intrusion of North Sea water into the Baltic occurred in 1975 (Tiews, 1976) following several years of increasing oxygen depletion in bottom waters. This situation of periodic stagnation, broken by saline inflows, followed by uplift of nutrients, favors periodic increase in biological production, unlike other areas of continuous anaerobiosis such as the deeper (below 100 m) zones of the Black Sea, which constitute a nutrient sink and are unproductive.

Oxygen depletion of bottom waters, with accompanying formation of hydrogen sulfide, occurred in Tokyo Bay in 1972 (Tsuiji et al., 1973; Seki et al., 1974), presumably related to an extensive red tide. Since red tides are becoming increasingly frequent in eutrophic bays in Japan as well as elsewhere in the world, anoxic conditions in bottom waters can be expected to increase in severity concomitantly. Mortalities of benthic organisms, associated with bottom water of low oxygen content, occurred in the Gulf of Trieste, in the North Adriatic in 1974 (Fedra et al., 1976). The authors reported scattered areas of decaying organisms in a region formerly characterized by stable benthic populations.

Oxygen depletion has occurred sporadically in Mobile Bay, Ala., one of the largest estuaries on the Gulf of Mexico. Stratification of the water column over highly organic bottom results in summer oxygen depletion, and occasionally, because of winds, the water mass impinges on beaches. Fish and invertebrates may be trapped in the anoxic water near beaches, often in a disoriented or moribund condition, where they are taken in great numbers by residents. The shoreline phenomenon is




Figure 6.—Distribution and relative abundance of *Ceratium tripos* bloom during early spring 1976.

called a "jubilee." Thirty-five such occurrences were reported by Loesch (1960) between 1946 and 1956, but newspaper accounts go back to the 19th century (the earliest being in 1867). May (1973) has reviewed the history of such events and finds no increase in their frequency in recent years. He carried out detailed oxygen determinations during a "jubilee" in 1971 and found large areas of the bay with less than 1 ppm dissolved oxygen in bottom water. Mortalities of fish, crabs, and oysters were observed.

There have been previous reports of

fish kills off New Jersey in 1968, 1971, and 1974 (Ogren, 1969; Ogren and Chess, 1969; Young, 1973, 1974²), and there may have been earlier problems that were not observed or reported. Those which were documented resemble the events of 1976 in that: 1) The more sedentary organisms found around rocks and wrecks and near open bottom were killed; 2) reports origi-

²Young, J. S. 1974. Unpublished data. Middle Atlantic Coastal Fisheries Center, National Marine Fisheries Service, NOAA, Highlands, N.J. Pers. commun. to J. B. Pearce.

nated from the same general area off central New Jersey; 3) depressed oxygen levels were considered a major contributing factor, and 4) suspended flocculent material was present in the water column. The 1976 episode differed from previous years in that: 1) It began earlier before the end of June, compared with the August-October period of earlier problems; 2) the bottom water was cooler (ca. 10°C) than reported in earlier publications (14°-18°C), and 3) hydrogen sulfide, not previously reported, was detected in lethal concentrations. It may have been present but not observed in earlier episodes.

The previous reports of mortalities also covered a much smaller area. The 1968 event, which appears to have been the most extensive of the earlier kills, included a zone from north of Long Branch to south of Barnegat Inlet, N.J. (a 70-km distance), from 1 to about 10 km offshore, with a total area of approximately 600-800 km² (Ogren and Chess, 1969). Mortalities were reported on and near wrecks and reefs from early September until late October 1968. Species affected were ocean pout (*Macrozoarces americanus*), cunner (*Tautoglabrus adspersus*), lobster (*Homarus americanus*), rock crabs (*Cancer irroratus*), mussels (*Mytilus edulis*), surf clams (*Spisula solidissima*), and starfish (*Asteria* sp.). More active species, such as tautog (*Tautoga onitis*), black seabass (*Centropristis striata*), squirrel hake (*Urophycis chuss*), conger eels (*Conger oceanicus*), and round scad (*Decapterus punctatus*), apparently were able to escape and were rarely reported among the mortalities. Fauna on wrecks off Barnegat and Atlantic City, N.J., was normal.

Reexamination of the same area in May and July 1969, disclosed that oxygen values near the bottom were in excess of 7.0 ml/l, and that wrecks had been repopulated by fish and crustaceans. No reports are available for 1970, but in early October 1971, lobsters and rock crabs were reported dead on several wrecks 12 km east of Manasquan Inlet in depths of about 30 m, and also at Shark River Inlet, north

of Manasquan Inlet (Young, 1973). Bottom water temperatures were high (18°C) and suspended flocculent material was noted low in the water column.

Again no reports are available for 1972 and 1973, but in August 1974, mortalities of ocean pout were observed on several wrecks off Manasquan Inlet (Young, footnote 2). Bottom dissolved oxygen values were 1.0 ml/l, with heavy suspended flocculent material and bottom water temperatures of 14°-15°C. In early September 1974, the *Subsea Journal* of the Manta Ray Diving Club of New Jersey reported dead lobsters and rock crabs on a wreck south of Manasquan Inlet.

Thomas et al. (1976) reported that significant summer depletion of bottom D.O. in the restricted area of the sludge and dredge spoil dump sites, and also in an area close to the New Jersey shore off Asbury Park, 10 km north of Manasquan Inlet, occurred during the summer of 1974. Low D.O. values have been reported previously in the New York Bight dump-site areas (Pearce, Segar and Berberian, 1976). D.O. values in dump-site areas in summer of 1975 were higher than 1974, and above the level considered harmful to most marine life.

CONCLUSION

The appearance and persistence of a massive bloom of the dinoflagellate *Ceratium tripos* in the Middle Atlantic Bight in 1976 is considered to be a significant factor contributing to oxygen depletion. Beginning in February, an unusual bloom of the dinoflagellate was found to be in progress over most of the outer continental shelf area of the Middle Atlantic Bight. This bloom was also noticed at many stations during an ichthyoplankton survey during March; the dinoflagellates clogged coarse-mesh nets from Long Island to Virginia in abundances which had not been observed in the 10 years previous experience of the ichthyoplankton group sampling in this area (Smith, footnote 1) (Fig. 6).

With onset of stratification, the bloom became concentrated in a band at or below the thermocline, within 100

km of the New Jersey coast. The *Ceratium* population appeared to decline sharply by July, and its contribution to bottom oxygen depletion in June and July was strongly indicated by the accumulation on the bottom of 1 cm or more of a brown flocculent material, which consisted to a large extent of dead *Ceratium*. Additionally, the observation in late spring and early summer that much of the bloom was concentrated at or below the thermocline indicates another possible mechanism of oxygen depletion, since the organism may exist heterotrophically. *Ceratium* is not grazed upon by many planktonic herbivores, so the persistent bloom over a period of several months meant the accumulation of large amounts of oxygen-demanding organic material in the water column, and the gradual accretion of dead organic material on the bottom. All of this oxygen-consuming material, when combined with the unusual hydrographic events mentioned earlier, seems to provide a reasonable explanation for the observed extreme oxygen depletion in 1976 (Mahoney, 1977; Malone, 1977).

Annual phytoplankton blooms are now a reality in sections of the New York Bight, probably in part at least as a consequence of organic loading of coastal waters, particularly from the Hudson River. The mean residence time calculated for the New York Bight is 100-250 days, and estimated daily chemical inputs from the Hudson River estuary complex alone are 520 t of nitrogen and 138 t of phosphate. Sewage sludge dumping and other human sources of nutrients have been estimated to augment these figures by only minor amounts (<10 percent).

In total though, when natural productivity of coastal waters is locally enriched by nutrients of human origin, a condition of increasing eutrophication can develop—and oxygen depletion can be a consequence. It does seem, however, that the extent and duration of the 1976 *Ceratium* bloom in the Middle Atlantic Bight would indicate that it did not occur in response to local nutrient inputs, but was rather a shelf-wide long-term phenomenon.

LITERATURE CITED

- Armstrong, R. S. 1977. Climatic conditions related to the occurrence of anoxia in the waters off New Jersey during the summer of 1976. In Oxygen depletion and associated environmental disturbances in the Middle Atlantic Bight in 1976, p. 17-35. National Marine Fisheries Service, Northeast Fisheries Center, Sandy Hook Laboratory, Tech. Ser. Rep. 3.
- Azarovitz, T. R., C. J. Byrne, and M. J. Silverman. 1977. A preliminary report recent and historic catches of finfish taken by ottertrawl in the vicinity of the 1976 New Jersey hypoxic water mass. In Oxygen depletion and associated environmental disturbances in the Middle Atlantic Bight in 1976, p. 365-415. National Marine Fisheries Service, Northeast Fisheries Center, Sandy Hook Laboratory, Tech. Ser. Rep. 3.
- Brongersma-Sanders, M. 1947. On the desirability of a research into certain phenomena in the region of upwelling water along the coast of South West Africa. Proc. K. Ned. Akad. Wet. 50(6):659-665.
- . 1957. Mass mortality in the sea. In J. W. Hedgepeth (editor), Treatise on marine ecology and paleoecology, Vol. 1, 941-1010. Copenhagen, W. J. 1953. The periodic mortality of fish in the Walvis region. A phenomenon within the Benguela current. S. Afr., Dep. Commer. Fish., Div. Fish., Invest. Rep. 14, 35 p.
- Davis, J. C. 1975. Minimal dissolved oxygen requirements of aquatic life with emphasis on Canadian species: a review. J. Fish. Res. Board Can. 32:2295-2332.
- Diaz, H. F. 1976. Meteorological conditions in the New York Bight during 1976. In Anoxia on the Middle Atlantic Shelf during the summer of 1976. National Science Foundation, Wash., D.C.
- Draxler, A., and C. J. Byrne. 1977. Hydrogen sulfide observations from R. V. *Albatross IV* and C.G.C. *Pt. Francis* cruises (26 Aug.-9 Sept., 21 Sept. 1976). In Oxygen depletion and associated environmental disturbances in the Middle Atlantic Bight in 1976, p. 83-91. National Marine Fisheries Service, Northeast Fisheries Center, Sandy Hook Laboratory, Tech. Ser. Rep. 3.
- Falke, H. Von. 1950. Das Fischsterben in der Bucht von Concepcion (Mittelchile). Senckenbergiana 31:57-77.
- Fedra, K., E. M. Ölscher, C. Scherübel, M. Stachowitsch, and R. S. Wurzian. 1976. On the ecology of a North Adriatic benthic community: distribution, standing crop and composition of the macrobenthos. Mar. Biol. (Berl.) 38:129-145.
- Festa, P. J. 1977. Observations on the summer flounder (*Paralichthys dentatus*) sport fishery in Great Bay, N.J. during the summer of 1976 in reference to anoxic water conditions. In Oxygen depletion and associated environmental disturbances in the Middle Atlantic Bight in 1976, p. 463-471. National Marine Fisheries Service, Northeast Fisheries Center, Sandy Hook Laboratory, Tech. Ser. Rep. 3.
- Freeman, B. L., and S. C. Turner. 1977a. The effects of anoxic water on the bluefish (*Pomatomus saltatrix*), an actively swimming fish. In Oxygen depletion and associated environmen-

- tal disturbances in the Middle Atlantic Bight in 1976, p. 431-450. National Marine Fisheries Service, Northeast Fisheries Center, Sandy Hook Laboratory, Tech. Ser. Rep. 3.
- _____, and _____. 1977b. The effects of anoxic water on the summer flounder (*Paralichthys dentatus*), a bottom-dwelling fish. In Oxygen depletion and associated environmental disturbances in the Middle Atlantic Bight in 1976, p. 451-462. National Marine Fisheries Service, Northeast Fisheries Center, Sandy Hook Laboratory, Tech. Ser. Rep. 3.
- Halgren, B. A. 1977. The effects of anoxic water conditions on the lobster fishery of New Jersey. In Oxygen depletion and associated environmental disturbances in the Middle Atlantic Bight in 1976, p. 417-429. National Marine Fisheries Service, Northeast Fisheries Center, Sandy Hook Laboratory, Tech. Ser. Rep. 3.
- Kalle, K. 1943. Die grosse Wasserumschichtung im Gotland-Tief vom Jahre 1933/34. Ann. Hydrogr. Berl. 71:142-146.
- Loesch, H. 1960. Sporadic mass shoreward migrations of demersal fish and crustaceans in Mobile Bay, Alabama. Ecology 41:292-298.
- MacKenzie, C. L., Jr. 1977. Effect of oxygen-deficient waters on the sea scallop (*Placopecten magellanicus*) off the New Jersey coast during summer, 1976. In Oxygen depletion and associated environmental disturbances in the Middle Atlantic Bight in 1976, p. 341-364. National Marine Fisheries Service, Northeast Fisheries Center, Sandy Hook Laboratory, Tech. Ser. Rep. 3.
- Mahoney, J. B. 1977. Observations on the decomposition of a *Ceratium tripos* population in New Jersey coastal waters during July - August 1976. In Oxygen depletion and associated environmental disturbances in the Middle Atlantic Bight in 1976, p. 233-251. National Marine Fisheries Service, Northeast Fisheries Center, Sandy Hook Laboratory, Tech. Ser. Rep. 3.
- Malone, T. C. 1977. The 1976 New Jersey fish kill: Summary of workshops. In Oxygen depletion and associated environmental disturbances in the Middle Atlantic Bight in 1976, p. ix-xxix. National Marine Fisheries Service, Northeast Fisheries Center, Sandy Hook Laboratory, Tech. Ser. Rep. 3.
- May, E. B. 1973. Extensive oxygen depletion in Mobile Bay, Alabama. Limnol. Oceanogr. 18:353-366.
- McHugh, J. L. 1977. Fisheries and Fishery Resources of New York Bight. U.S. Dep. Commer., NOAA Tech. Rep., NMFS Circ. 401, 50 p.
- Meyer, P. F., and K. Kalle. 1950. Die Biologische Umstimmung in der Ostsee in den lentzen Jahrzehnten, eine Folge hydrographischer Wasserumschichtungen. Arch. Fischereiwiss. 2:1-9.
- National Marine Fisheries Service. 1977. Oxygen depletion and associated environmental disturbances in the Middle Atlantic Bight in 1976. National Marine Fisheries Service, Northeast Fisheries Center, Sandy Hook Laboratory, Tech. Ser. Rep. 3, 471 p.
- National Marine Fisheries Service. 1973. Sportfish emphasis document, Middle Atlantic Coastal Fisheries Center. National Marine Fisheries Service, Middle Atlantic Coastal Fisheries Center, Informal Rep. 15, 79 p.
- Ogren, L. 1969. New Jersey fish/crustacean mortality. Smithsonian Institution, Center for Short Lived Phenomena, Event No. 38-69, Card No. 509, 510.
- Ogren, L., and J. Chess. 1969. A marine kill on New Jersey wrecks. Underwater Nat. 6(2):4-12.
- Pearce, J. B. 1972. The effects of solid waste disposal on benthic communities in the New York Bight. In M. Ruivo (editor), Marine pollution and sea life, p. 404-411. Fishing News (Books) Ltd., Lond.
- Radosh, D. J., A. Frame, and T. Wilhelm. 1977. Short-term effects on the benthic infauna resulting from oxygen-deficient bottom waters off the New Jersey coast, 1976; preliminary report. In Oxygen depletion and associated environmental disturbances in the Middle Atlantic Bight in 1976, p. 109-126. National Marine Fisheries Service, Northeast Fisheries Center, Sandy Hook Laboratory, Tech. Ser. Rep. 3.
- Ropes, J. W., and S. Chang. 1977. Impact on offshore sea clam populations (surf clam and ocean quahog) associated with anoxic bottom water in the Middle Atlantic Bight during summer, 1976. In Oxygen depletion and associated environmental disturbances in the Middle Atlantic Bight in 1976, p. 263-340. National Marine Fisheries Service, Northeast Fisheries Center, Sandy Hook Laboratory, Tech. Ser. Rep. 3.
- Segar, D. A., and G. A. Berberian. 1976. Oxygen depletion in the New York Bight apex: Causes and consequences. Am. Soc. Limnol. Oceanogr., Spec. Symp. 2:220-239.
- Seegerstråle, S. G. 1969. Biological fluctuations in the Baltic Sea. Progr. Oceanogr. 5:169-184.
- Seki, H., T. Tsuji, and A. Hattori. 1974. Effect of zooplankton grazing on the formation of the anoxic layer in Tokyo Bay. Estuarine Coastal Mar. Sci. 2:145-151.
- Sindermann, C. J. 1970. Principal diseases of marine fish and shellfish. Acad. Press, N.Y., 369 p.
- _____. 1976. Oyster mortalities and their control. FAO technical conference on aquaculture, Kyoto, Japan. DOC FIR: Aq.76/R34, 25 p.
- Steimle, F. 1977a. The persistence and boundaries of the bottom water oxygen depletion problem of 1976 in the New York Bight. In Oxygen depletion and associated environmental disturbances in the Middle Atlantic Bight in 1976, p. 41-64. National Marine Fisheries Service, Northeast Fisheries Center, Sandy Hook Laboratory, Tech. Ser. Rep. 3.
- _____. 1977b. A preliminary assessment of the impact of the 1976 N. Y. Bight oxygen depletion phenomenon on the benthic invertebrate megafauna. In Oxygen depletion and associated environmental disturbances in the Middle Atlantic Bight in 1976, p. 127-165. National Marine Fisheries Service, Northeast Fisheries Center, Sandy Hook Laboratory, Tech. Ser. Rep. 3.
- Theede, H., A. Ponat, K. Hiroki, and C. Schlieper. 1969. Studies on the resistance of marine bottom invertebrates to oxygen deficiency and hydrogen sulfide. Mar. Biol. (Berl.) 2:325-337.
- Tiews, K. 1976. Ostsee: GröÙe Wassererneuerung seit Jahrzehnten. Umsch. Wiss. Tech. 76:271.
- Thomas, J. P., W. C. Phoel, F. W. Steimle, J. E. O'Reilly, and C. A. Evans. 1976. Seabed oxygen consumption - New York Bight, U.S.A. Am. Soc. Limnol. Oceanogr., Spec. Symp. 2:354-369.
- Tsuji, T., H. Seki, and A. Hattori. 1974. Results of red tide formation in Tokyo Bay. J. Water Pollut. Control Fed. 46:165-172.
- Young, J. S. 1973. A marine kill in New Jersey coastal waters. Mar. Pollut. Bull. 4:70.

MFR Paper 1356. From Marine Fisheries Review, Vol. 40, No. 12, December 1978. Copies of this paper, in limited numbers, are available from D822, User Services Branch, Environmental Science Information Center, NOAA, Rockville, MD 20852. Copies of Marine Fisheries Review are available from the Superintendent of Documents, U.S. Government Printing Office, Washington, DC 20402 for \$1.10 each.

Townsend Cromwell Charts Guam, Marianas Marine Resources

The NOAA ship *Townsend Cromwell* operated by the National Ocean Survey and attached to the National Marine Fisheries Service in Honolulu, returned to Honolulu on 21 July after completing a 10-week survey and assessment cruise to waters around Guam and the Northern Mariana Islands. The primary mission for the cruise was to investigate the marine resources around these islands and identify those that have the greatest potential for early development. According to Richard S. Shomura, Director, Honolulu Laboratory, National Marine Fisheries Service, NOAA, this cruise was a joint effort of NMFS, the Guam Division of Fish and Wildlife, the University of Guam Marine Laboratory, and the Northern Marianas Fishing Authority. This timed major survey of the waters around Guam and the Northern Marianas will provide information which the governments of Guam and

the Northern Marianas can use in planning their fishery development programs, said Shomura.

Using a variety of fishing gear including traps, handline, bottom trawls, and trolling lines, scientists aboard the *Townsend Cromwell* found relatively few areas with high densities of commercially valuable species such as ulua or jacks, snappers, and pandalid shrimps. However, bigeye scad or akule was notably abundant along the seamounts such as Arakane Reef, Pathfinder Reef, and other unnamed seamounts. Night "jigging" for bigeye scad during the dark moon phases of the month was particularly effective at some of the night-light fishing stations over the seamounts, said Richard N. Uchida, Chief Scientist on the cruise. Handline fishing produced good catches of gindai and ulua at several of the offshore seamounts.

The survey also revealed that rela-


tively high densities of pandalid shrimps occur in certain areas around Guam and the Northern Marianas. According to Uchida, these shrimps were caught in pots set in waters ranging from 365 to 820 m in depth. The catch rate was highest in waters off Guguan Island where it reached 175 shrimps per pot at depths of 640-730 m.

Trolling was also very productive over some of the offshore seamounts and catches included yellowfin tuna, skipjack tuna, and rainbow runner. Of 226 fish caught by trolling, 114 were tagged and released. When recaptured, these tagged fish will provide information on migration patterns and growth.

In general, it appeared that the offshore seamounts were more productive fishing grounds compared with locations close to the high islands, said Uchida. The handlining catch reached 1.9 fish per line-hour over the seamounts, whereas it was only 1.0 fish per line-hour at stations near the high islands. Of course, these catch rates are based only on the results of one cruise and may be due to seasonal variation in the distribution and abundance of the fish, cautioned Uchida. It was also noted that two species of spiny lobsters are known to occur in the Northern Marianas and both were taken in lobster traps during the cruise. However, these species are primarily reef-dwelling forms and very few were caught in the offshore zone where the *Townsend Cromwell* fished. Likewise, trawling for bottom fish using a Norwegian fish trawl was generally unproductive. The *Townsend Cromwell* also conducted coral-drag operations in depths ranging from 42 to 450 m of water, but no potentially valuable precious coral grounds were located.

COMPUTERS MAY AID FISHING INDUSTRY

Can computers reduce the risk of failure in the fishing business? Economists at five universities across the country, supported by a \$138,100 grant from NOAA, think so and are setting out to prove it.



Ned A. Ostenso, Director, Office of Sea Grant, said aquaculture and fishing ventures traditionally have been subject to financial loss or failure because of unexpected rises in costs, and other problems. "This project aims to develop a computer-model budget and the necessary data base to help the fishing industry improve its chances of success," he said.

The project will bring together experienced economists from Texas A&M University, the University of Massachusetts, Oregon State University, Louisiana State University, and the University of Rhode Island to develop necessary analytical tools and techniques that others can use.

A general computer program will be developed that can be applied to most aquaculture systems and many different fishing vessels to predict the probable outcome of a particular venture. Economic analyses then will be prepared for high priority species of fish within each region of the country represented by the participating scientists.

Data for the aquaculture phase of the project will include such species as salmon, marine shrimp, oysters, freshwater prawns, and lobster. Information on vessels which fish specifically for lobster, groundfish, shrimp, snapper-grouper, tuna, crabs, salmon, and halibut also will be included. Problems will range from determining which type of equipment would be most cost effective for a particular aquaculture venture to assessing the effect of changes in tax laws for vessel owners with respect to crews and imported products used in fishing. The five cooperating universities will provide \$82,400 in matching funds.

LSU DESIGNATED A SEA GRANT COLLEGE

Louisiana State University (LSU) has been designated a Sea Grant College, Secretary of Commerce Juanita M. Kreps has announced, in recognition of the excellence of the institution's marine resources program. LSU is the 13th institution in the United

States to attain Sea Grant College status.

"The Sea Grant program at Louisiana State University, now in its eleventh year, has served as the stimulus for development of a broad program in marine affairs," Secretary Kreps said. "The focus of education, research, and marine advisory service components of the program is on the wise use, management, and conservation of renewable natural resources offshore and in the coastal zone."

Ned A. Ostenso, Director of the National Sea Grant Program, an element of the National Oceanic and Atmospheric Administration (NOAA), said the designation underlines the institution's strong development in marine resource activities and the cooperation between the University, the State, and the Federal Government. The National Sea Grant Program has awarded grants to universities, laboratories, and other institutions since 1968, supporting research, education, and advisory services that foster development of the nation's marine resources. Operating under a matching fund arrangement, the Sea Grant Program includes more than 690 projects at approximately 130 colleges, universities, and institutions.

The Sea Grant Program at LSU is centered on the University's Baton Rouge campus, where, according to LSU System President Martin D. Woodin, it has had a deep influence resulting in a strong commitment to marine affairs. "An example of this has been the establishment of the Center for Wetland Resources, which includes the Coastal Studies Institute, the Laboratory for Wetland Soils and Sediments, the Coastal Ecology Laboratory, and the Department of Marine Sciences," he said.

Sea Grant research activities at LSU have focused to a large extent on local and regional needs. Detailed ecological studies were made in the Barataria Basin, for example, which led to major findings regarding productivity of the coastal marshes and related effects of human activity. In other areas, LSU Sea Grant researchers have achieved success in studies dealing with international and coastal law, in projects on

rearing of shrimp and crawfish, in a program of immunization for farm-raised alligators, in expansion of public understanding of the oceans at all age levels, and in a marine advisory service program designed to uncover problems faced by fishermen, aquaculturists, food processors, and other marine users for presentation to researchers and for relaying research results to those users.

SATELLITE CHECKS OCEAN POLLUTION

Two highly complex instruments carried into space with the launch of the Nimbus-G satellite in late 1978 hold promise of providing scientists with the answers to two basic questions of importance to mankind: 1) How polluted are the world's oceans becoming, and 2) is the earth warming up or cooling down? The instruments are vital to studies being conducted by scientists with the National Oceanic and Atmospheric Administration (NOAA), as well as to other researchers concerned with the oceans and the atmosphere.

Nimbus-G, a research and development satellite managed for the National Aeronautics and Space Administration by its Goddard Space Flight Center, Greenbelt, Md., was scheduled for launch from NASA's Western Test Range in Lompoc, Calif. One instrument, the Coastal Zone Color Scanner, is expected to aid oceanographers in determining the content of water, important in monitoring water pollution, according to Warren A. Hovis, Jr., Director of the Satellite Experiment Laboratory of NOAA's National Environmental Satellite Service. NOAA is a Commerce Department agency.

The scanner, sensing the colors in water beneath the polar-orbiting satellite, will permit content analyses to be made of large areas of coastal or ocean waters, "letting oceanographers view the ocean as never seen from ships," Hovis said. The instrument will be used to determine how well water pollution—such as oil spills, sewage and industrial waste dumpings, and river sediment—can be detected and tracked.

The other instrument, the Earth

Radiation Budget experiment, makes a variety of measurements of the radiation coming from the sun and the earth. Of particular interest to scientists is the observation of variations in the radiative heat exchange between the sun and earth with time, with location on the earth, and over the entire globe.

NOAA's Herbert Jacobowitz, of the Satellite Service's Meteorological Satellite Laboratory, said such observations can help in monitoring, and perhaps eventually anticipating, the fluctuations in climate from months to a few years. This, he explained, is because this radiative heat exchange creates the basic energy source for the atmospheric and oceanic circulations that determine climate. For example, variations in radiative heating between the tropics and higher latitudes, or between the oceans and continents, may yield important clues as to the types of winters or summers that may be expected to occur in the United States or other parts of the world.

New Technique Tracks Sewage Dumped At Sea

National Oceanic and Atmospheric Administration (NOAA) scientists have identified a biochemical technique that can be used to trace sewage in coastal areas, a matter of vital interest to environmental managers in heavily populated parts of the coast. The newly identified technique uses coprostanol, a steroid thought to be produced exclusively by bacteria in the intestines of mammals, to measure sewage in offshore sediments.

According to Patrick G. Hatcher, Philip A. McGillivray, and NOAA Corps Lieutenant Commander Larry E. Leister, all of NOAA's Atlantic Oceanographic and Meteorological Laboratories in Miami, the coprostanol method promises to become the standard for sewage-pollution detection. Hatcher and his colleagues developed the technique as part of a major ecosystem study NOAA has conducted in the New York Bight, the 15,000-square-mile (39,000-km²) continental shelf area off the New York-New Jersey coast.

Because sewage sludge, the end product of waste-water treatment, is dumped in the Bight, scientists and citizens are concerned about sewage contamination there. But sewage is difficult to trace in the chemically complex marine environment, Hatcher said. The Commerce Department researchers turned to steroids, biochemical compounds that resist deterioration in the environment. The steroid coprostanol was found to be present in contaminated marine sediments, but not in uncontaminated sediments, and appeared to be stable enough to be used as a tracer of sewage. Subsequent work led to a "percent-coprostanol" term that permits the scientists to estimate how much of a sample's organic matter is sewage-derived.

With the technique, a map of the New York Bight has been developed showing sewage pollution centered in a basin near the sewage-sludge dump site, and diminishing rapidly with distance away from the dump site. The highest value found was 15 percent coprostanol in the highly contaminated black muds near the dump site. Pure sewage contains more than 30 percent.

Substances From Sea Creatures May Have Medical Applications

Twelve chemical compounds recently discovered in such sea creatures as sponges, sea cucumbers, and sea hares potentially are valuable in treating cancer, central nervous system disorders, and cardiovascular problems, researchers supported by the National Oceanic and Atmospheric Administration (NOAA) believe. The scientists, from the University of Oklahoma, have isolated and identified the compounds during the past 15 months, and now are beginning further study on their application to disease. The research is supported by a \$116,500 Sea Grant from NOAA, a Commerce Department agency, and \$58,258 in supplemental funds from the University.

One of the most promising substances for the treatment of nervous


disorders is found in the sea hare—a member of the shellfish family—and is known as dactylene. The substance may inhibit the breakdown in the body of barbiturates, which are used to induce sleep and can be addictive. Inhibition of their breakdown would prolong their effects and permit use of smaller and safer doses. Dactylene has been tested on rats and mice, and increases the length of time the laboratory animals sleep after being given a barbiturate.

Other newly identified compounds suppress cellular growth in tissue and are candidates for possible anticancer treatment, according to the research team. The scientists are now conducting experiments on laboratory animals to validate the effects of the substances.

Chemicals taken from a number of Caribbean invertebrates and algae have, in the laboratory, inhibited the growth of experimental tumors in mice, and also mitigated cardiovascular and central nervous system disorders induced in small mammals. Isolation and identification of these substances will continue under the NOAA grant.

The newly discovered compounds may serve as models for the synthesis of new drugs. Additionally, the chemical information gained from the research will increase knowledge of the chemistry of marine animals and plants, and define the types of compounds released into seawater from natural sources.

Elephant's ear sponge.



Peru Reduction Fishery Drops 50 Percent in 1977

Peru's 1977 catch of fish for reduction decreased substantially to 2,020,000 metric tons (t), or 50 percent less than in 1976. The country's total fisheries catch in 1976 was 4,343 million metric tons, according to FAO statistics. While the country was able to get through the year without major changes in its imports, prospects for an even lower 1978 fisheries catch were expected to result in considerably larger soybean oil imports.

Table 1.—Peru's fish oil supplies and distribution in calendar years 1976-78. Source: Ministry of Fisheries, Trade and Attache's estimates.

Item	1976 (Final)	1977 (Preliminary)	1978 (Forecast)
Supplies (x 1,000 t)			
Opening stocks (1 Jan.)	7	8	16
Production	104	101	160
Total	111	109	76
Distribution			
Exports	— ²	0	0
Apparent consumption			
Edible	98	83	56
Industrial ³	5	10	10
Subtotal	103	93	66
Ending stocks (31 Dec.)	8	16	10
Total distribution	111	109	76

¹Based on a 1.5 million t fish catch (anchovy, sardines, jurel, and needlefish).

²20 t hydrogenated fish oil exported to Bolivia.

³Soap industry.

Table 2.—Peru's consumption of fats and oils in calendar years 1976-78. Sources: Trade and Attache's estimates.

Item	1976 (Final)	1977 (Preliminary)	1978 (Forecast)
Fish oil (x 1,000 t ¹)			
Liquid oil ²	45.1	43.7	16.0
Solid fats (margarine and shortening)	53.3	38.8	40.0
Soap industry	5.0	10.0	10.0
Total	103.4	92.5	66.0
Per capita consumption, edible, (kg/year)			
Vegetable oil	6.1	5.7	7.3
Fish oil	6.2	5.0	3.3
Lard	0.5	0.5	0.5
Total	12.8	11.2	11.1

¹Unless otherwise indicated.

²For making blended cooking oil.

Peru's production of fats and oils was also forecast to decrease from 135,300 t in 1977 to only 97,200 t in 1978, down 28 percent, due to the unfavorable anchovy fishery. The output of fish oil was forecast to decline 40 percent in 1978 (Table 1).

In a surprising development, the Ministry of Fisheries authorized the catch of sardines, jurel, and needlefish as of 17 April 1978. Publicly announced forecasts estimated the 1978 catch of reduction fish at 1.5 to 1.8 million tons, fish meal production at 300,000 to 350,000 tons, and fish oil production at 60,000 to 70,000 tons. All figures are revised slightly downward from earlier estimates.

In 1976 and 1977, fish oil represented roughly 46 percent of Peru's total consumption of fats and oils. Based on the 1978 unfavorable fishing forecasts, fish oil should constitute only about 32 percent of Peru's total fats and oils requirement this year. According to various sources, no more than 120,000 t of fish oil (4 million t anchovy catch) is expected in the coming years.

Tables 2 and 3 show the supply and distribution of fish oil and fish meal, respectively, for calendar years 1976-78. As can be seen by Table 3, a reduction in fish meal consumption of 35,000

Table 3.—Peru's fish meal supplies and distribution in calendar years 1976-78. Sources: Ministry of Fisheries, Trade and Attache's estimates.

Item	1976 (Final)	1977 (Preliminary)	1978 (Forecast)
Supplies (x 1,000 t)			
Opening stocks (1 Jan.)	40	172	112
Production	886	493	300
Total	926	665	412
Distribution			
Exports	624	436	300
Apparent consumption	130	117	82
Ending stocks (31 Dec.)	172	112	30
Total	926	665	412

¹Based on a 1.5 million t fish catch (anchovy, sardines, jurel, and needlefish).

t from 1977 to 1978 was expected. The decline in 1978 fish oil and fish meal production is due to the reduced fisheries catch. Domestic production has a direct bearing on the availability of fish meal for export. (Source: IFR-78/102.)

Japan, Russia Negotiate Joint Fishing Ventures

Soviet Fisheries Minister Aleksandr A. Ishkov informed Japan Fisheries Association President Tomoyoshi Kamenaga earlier this year that his Ministry had received five or six applications from Japanese trading firms to form joint ventures with the Soviet Union. The Japanese companies were seeking access to the Soviet 200-mile zone in areas now closed to Japanese fishing vessels under the existing Provisional Japan-Soviet Fisheries Agreement.

The two men met the day after the signing of the Five-Year Japan-Soviet Fisheries Cooperation Agreement in Moscow. Ishkov stated that the U.S.S.R. was willing to enter into several joint ventures as soon as the Japanese Government determined which requests would be approved. The deadline set for the Japanese response was 10 May. Conditions set forth then by Ishkov were: 1) Fishing vessels of the two countries would operate separately in the same area; 2) 25 percent of the Japanese catch would be transferred to the U.S.S.R. as a fishing fee; and 3) Japan would buy back this 25 percent as well as purchase the Soviet catch.

The Japanese Fisheries Agency (JFA) had approved, as of 6 June, five of the six proposed joint operations. The JFA indicated that it would only

Unless otherwise noted, material in this section is from the Foreign Fishery Information Releases (FFIR) compiled by Suneo C. Sonu, Foreign Reporting Branch, Fishery Development Division, Southwest Region, National Marine Fisheries Service, NOAA, Terminal Island, CA 90731, or the International Fishery Releases (IFR) or Language Services Daily (LSD) reports produced by the Office of International Fisheries, National Marine Fisheries Service, NOAA, Washington, DC 20235.

permit those Japanese companies to participate which previously fished within the Soviet 200-mile zone. All Japanese trading companies would be excluded. This decision by the JFA was delayed past the 10 May deadline proposed by Ishkov owing to the following two concerns.

1) The joint operation plans should be conducted separate from the Japan-Soviet Provisional Fisheries Agreement and should not adversely affect the 850,000 metric ton (t) 1978 Soviet quota for Japanese vessels fishing within the Soviet 200-mile zone.

2) Japanese fishing companies and organizations should take the initiative in concluding contracts with the U.S.S.R., although the trading companies originated the plans and filed the applications.

According to later information received from the U.S. Regional Fisheries Attache in Tokyo, the JFA and the Soviets agreed to proceed with four of the six proposed joint ventures (Table 1). Alaska pollock was excluded because it is a migratory fish and not limited to the continental shelf. Blue king crab was also deleted, apparently at the insistence of the Soviets (Table 2).

The Japanese have agreed to provide the Soviets with information on Japanese fishing methods, especially basket-net fishing, and the Soviets will place fishermen as observers on Japanese vessels involved in the joint operations. The question of fees is still being negotiated. The Japanese Government will decide about participation in joint ventures next year after viewing this year's results. Japanese vessels will be issued temporary licenses and will not be eligible for government compensation should the program be discontinued next year.

According to the NMFS Office of International Fisheries the status of the proposed Soviet-Japanese joint ventures at midsummer was unclear. The Japanese Government announced on 22 June that they had recently been informed by the Soviet Union that the Soviets would not be able to carry out the joint fishing ventures as had been agreed. The reason given, according to a Japanese Government source, was

Table 1.—U.S.S.R.-Japan planned joint fishing operations, 1978.

Japanese company or organization		Species	Quota (t)	Area	Vessels					
					Japanese			Soviet		
Fishery	Trading				No.	Type	Size ¹	No.	Type	Size ¹
Hoko Suisan & Nichiro Gyogyo	Marubeni	Tanner crab	3,000	Olyutorskii Bay	1	Reefer	1,000	2	NA	150
					2	Sub-reefers	400			
					3	Pot-vessels	96			
Hamaya Suisan ²	Tokyo Maruichi Shoji	Pink shrimp	700	Southwest Kamchatka	1	Shrimp trawler	NA	1	NA	NA
Japan Sea Shrimp Pot Fishing Council	Yokohama Tsusho	Pink shrimp	700	Marmiya Straits	7	Pot-vessels	96	2	NA	NA
Okhotsk Sea Hair Crab Fishing Council	Tokyo Maruichi Shoji	Hair crab	500	Southwest Sakhalin	1	NA	299	1	NA	NA
					1	NA	NA			

¹Gross registered ton per vessel.

²The company is located in Wakkanai, Hokkaido.

Source: Regional Fisheries Attache, U.S. Embassy, Tokyo.

Table 2.—U.S.S.R.-Japan joint fishing operations not approved, 1978.

Japanese company or organization		Species	Quota (t)	Area	Vessels					
					Japanese			Soviet		
Fishery	Trading				No.	Type	Size ¹	No.	Type	Size ¹
Medium Trawler Owners	Tairiku Boeki	Pollock	20,000	Southern Kamchatka	12	Medium trawlers	NA	— None —		
Taiyo Gyogyo	Morikawa Shoji	Blue king crab	1,000	Olyutorskii, Naarin Bays		NA	400	— None —		

¹Gross registered ton per vessel.

Source: Regional Fisheries Attache, U.S. Embassy, Tokyo.

Moscow's displeasure with the resumption of treaty talks between Japan and the People's Republic of China. This announcement came just as the joint ventures were about to begin operations. It was also announced at that time, that the Soviets would not allow the resumption of kelp (konbu) gathering by small-scale Japanese fishermen off Soviet-held Kaigara Island.

An official Moscow source denied in

23 June that the Soviets had cancelled the joint ventures. At the same time, this official source would not say definitely whether negotiations on the joint ventures and kelp gathering would continue as before. Even so, on 4 July, the Soviet Union notified the Japan Fisheries Association that there would be a further delay in joint venture talks owing to disagreements among Soviet authorities. (Source: IFR-78/125.)

Kamaboko Reportedly Made From Krill

Japan's National Federation of Kneaded Fisheries Products Cooperatives announced last summer the successful development of "kamaboko", a heat-pasteurized fishcake, using Antarctic krill. The test product reportedly excelled the quality of the Alaska pollock-based kamaboko in nutritional value and taste, and possessed desirable flavor, color, and texture.

Commercial production was expected to encounter such problems as difficulty of using ordinary mechanical peelers, low yield of krill meat (averaging about 8 percent of body weight), and a strong tendency for the krill meat to oxidize, which would necessitate mothership production of krill-based kamaboko.

(Source: FFIR 78-10.)

NMFS Scientific Reports Published

NOAA Technical Report NMFS SSRF-720. Squire, James L., Jr. **"Sea surface temperature distributions obtained off San Diego, California, using an airborne infrared radiometer."** March 1978. 30 p. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, DC 20402.

ABSTRACT

Sea surface temperature surveys were conducted weekly off San Diego, Calif., using an airborne infrared radiometer from April through October 1972-74. A total of 90 surveys were made over the 320-mile flight track. The analog chart record of temperature was keyed to a "ground truth" temperature measurement and read to determine 1 min average temperatures which were plotted on the flight track and 1°F (0.56°C) isotherms were contoured from the data. The trend of sea surface temperatures during the 3-year period showed warmer temperatures in 1972, which was an "el Niño" year, cooler in 1973, warmer than 1973 in 1974, but not as warm as 1972. In early July 1974, an anomalous warming period occurred and highest average temperatures of 73°F (22.7°C) were recorded in 19 July 1974. Lowest average temperature of 54°F (12.2°C) was recorded on 2 April 1973.

The 1972-74 survey temperatures taken over the "ground truth" calibration site were compared with a time series of temperature observations taken during the same months from 1963 to 1968. The average temperature during 1963-68 was 63.4°F (17.4°C) and for 1972-74,

64.1°F (17.8°C), a difference of less than 1°F (0.56°C). Average monthly temperature differences, 1963-68 compared with 1972-74, shows April the same, warmer 2°F (1.12°C) for May, 3°F (1.68°C) for June, 1°F (0.56°C) for July, same for August, 1°F (0.56°C) less for September, and 2°F (1.12°C) less for October.

NOAA Technical Report NMFS SSRF-725. Davis, Clarence W. **"Seasonal bottom-water temperature trends in the Gulf of Maine and on Georges Bank, 1963-75."** May 1978. 17 p.

ABSTRACT

Spring (1968-75) and autumn (1963-75) bottom-water temperature temperatures in the Gulf of Maine and on Georges Bank were analyzed to investigate a suspected warming trend in the region. During the spring the mean temperature in the Gulf of Maine increased rather steadily from a low of 5.4°C in 1968 to a high of 6.4°C in 1974. Various subareas of the Gulf had more frequent and greater oscillations but exhibited the same overall warming trend. Mean spring temperatures on Georges Bank fluctuated from 3.8°C in 1968 to 6.3°C in 1974 and declined by nearly 2°C in 1975 with similar characteristics in eastern, central, and western subareas of the Bank.

During the autumn in the Gulf, bottom-water temperatures reached a minimum of 5.4°C in 1966, increased to a maximum of 8.4°C in 1973 and 1974, but declined to 8.0°C in 1975. Subareas of the Gulf

generally showed the same temperature trends from 1963 to 1968; especially notable are the cooling trend west of long. 69°W which commenced in 1971, and a decrease in all five subareas in 1975. Georges Bank temperatures in autumn declined from a maximum of 13.1°C in 1965 to a minimum of 10.4°C in 1969, reached another peak of 12.6°C in both 1973 and 1974, but declined to 11.6°C in 1975. Subareas of Georges Bank generally followed the same pattern with the eastern third of the Bank usually 2°C or colder than either of the other subareas in the autumn.

The average bottom-water temperatures during spring were 5.0°C on Georges Bank and 6.1°C in the Gulf of Maine; temperatures in the autumn were 11.7°C and 7.2°C, respectively, for these areas.

NOAA Technical Report NMFS Circular 411. Dooley, James K. **"Systematics and biology of the tilefishes (Perciformes: Branchiostegidae and Malacanthidae), with descriptions of two new species."** April 1978. 78 p.

ABSTRACT

Tilefishes have been examined on a world basis with the following conclusions: 1) Tilefishes belong to two distinct phyletic lines here designated as the family Branchiostegidae and the resurrected family Malacanthidae. 2) The Branchiostegidae include 3 genera and 21 species. 3) The Malacanthidae include 2 genera, 2 subgenera, and 8 species. 4) A new species, *Caulolatilus hubbsi*, is described from off southern California, the Gulf of California, the Galapagos Islands, and from Callao, Peru; it is generally found sympatrically with the other two eastern Pacific species of *Caulolatilus*. 5) A second new species, *Branchiostegus albus*, is described from off central Honshu, Japan; Pusan, Korea; along the coast of the East China Sea including Shanghai and Taiwan; and the coasts of the South China Sea including Hong Kong and Macao. *Branchiostegus albus* was formerly confused with *B. argentatus* Cuvier 1830. 6) Branchiostegids are generally relatively deep dwelling (20-600 m;

usually deeper than 50 m) fishes found along the edges of continental margins, near the upper slope of islands, or at the heads of deep-sea canyons. These fishes are deep bodied and have prominent skull crests. 7) Malacanthids are relatively shallow-water (10-150 m, usually shallower than 50 m), burrow-dwelling or mound-building fishes with elongate bodies and rounded or flat skulls with no prominent crests. 8) Tilefishes appear to have basal percoid affinities, having a number of larval and osteological characters found among beryciform fishes (considered antecedent to perciforms) and characters considered primitive among perciform fishes.

NOAA Technical Report NMFS SSRF-721. Hall, R. A., E. G. Zook, and G. M. Meaburn. "National Marine Fisheries Service survey of trace elements in the fishery resource." March 1978. 313 p.

ABSTRACT

Trace element levels have been determined in tissues of 204 species of finfish, Mollusca, and Crustacea taken from 198 sites around the coastal United States, including Alaska and Hawaii. The survey was undertaken as part of the Microconstituents Program of the National Marine Fisheries Service, and covers the occurrence of 15 elements: antimony, arsenic, cadmium, chromium, copper, lead, manganese, mercury, molybdenum, nickel, selenium, silver, tin, vanadium, and zinc. Total concentrations of each element were determined without regard to chemical form. The species analyzed represent approximately 93 percent of the volume of the U.S. commercial and sportfish catch. The analytical data are summarized in several ways in order to emphasize different aspects of the trace element distributions. Mean levels of each element are presented in relation to the number of species examined, the U.S. (commercial and sportfish) catch, and the U.S. catch intended for consumption. More detailed analytical data on all 15 elements are given for individual species with reference to tissue analyzed, length and weight of fish,

and location of catch. For the most part, experimental results are presented without interpretive comment. Mean levels of mercury, the only element for which a regulatory

action level is in force, were found to exceed 0.5 ppm Hg in species representing less than 2 percent of the U.S. catch intended for consumption.

Marine Recreational Fishing Symposium Proceedings Printed

Marine Recreational Fisheries 2 contains the proceedings of the Second Annual Marine Recreational Fisheries Symposium and was published by the Sport Fishing Institute in Washington, D.C. Objectives of the symposia series are to identify major recreational marine fisheries problems—biological, economic, and social—and to promote effective management practices, based on scientific principles, for the conservation of living marine resources.

The book's 20 chapters (symposium presentations plus discussions) were delivered by leading national and international authorities, many with the National Marine Fisheries Service. A talk on NMFS mission and goals and the 200-mile zone law was given by then-NMFS Director Robert Schoning. Richard Hennemuth contributed a talk on "Some Biological Aspects of Optimum Yield"; Grant Beardsley and Wesley Parks presented "Management of Western North Atlantic Bluefin Tuna Fisheries"; and Brian Rothschild, J. M. Gates, and A. M. Carlson coauthored "Management of Marine Recreational Fisheries." "Protection of

Coastal Wetlands" was explained by Richard Gardner, acting director of NOAA's Office of Coastal Zone Management.

Other chapters discuss marine angling perspectives in both Canada and Mexico, the U.S. 200-mile zone law, criteria for collecting marine recreational fisheries data, attaining "clean water", effects of water development on striped bass, the tarpon's unusual biology and man's impact on its future, economic losses to marine recreational fisheries from habitat destruction, a model for determining optimum yield, human perspectives in optimum sustainable yield fisheries management, and "menhaden, sport fish and fishermen."

Symposium chairman was Richard H. Stroud of the Sport Fishing Institute; proceedings editor was Henry Clepper. The 220-page hardbound volume is available at \$15 per copy from the International Game Fish Association, 3000 E. Las Olas Blvd., Fort Lauderdale, FL 33316. It will be a valuable addition to the marine recreational angling literature.

NEKTON VOLUME IS TRANSLATED

Nekton, by Yu. G. Aleyev, has been translated from the Russian by B. M. Meerovich and published by Dr. W. Junk b.v. - Publishers, P.O. Box 13713, 2501 ES The Hague, The Netherlands. The book is divided into three parts: 1) Systematics and geographical distribution of nekton; 2) fundamental nektonic adaptations; and 3) origins and ecological divergence of nekton. It provides a good review of Russian studies on the topic.

In the introduction, the author defines his terms, presents the history and

task of nektonology, and describes nektonological investigation methods. Other chapters discuss maintaining the body suspended in the water, locomotion, reducing resistance to movement, controlling movement, camouflage and defense, other adaptations, classes of nekton and their origins, and nekton and the body of water. The 435-page book has 247 figures, contains an extensive literature cited section, indexes of author's names and animal latin names, plus a subject index. The listed price is 120.00 Dutch guilders.

Authors, Titles, and Subjects in *Marine Fisheries Review* Volume 40, Numbers 1-12, 1978

Indexed here by author, title, and subject are the papers which appeared in *Marine Fisheries Review* in 1978. A list of these papers, in numerical order, appears at the end of the Index. Entries are indexed by number and page (i.e., 7:12 indicates the July number, page 12).

Anonymous notes, news articles, and regular departments are not indexed.

A

- Aequipecten irradians*—see Scallop, bay
Aeromonas salmonicida
bacterins
used in fish immunization, 3:12
Alaska, southeastern
herring roe gillnet fishery, 1976
economic aspect of development, 4:25
Alosa sapidissima—see Shad, American
“(An) analysis of the charter boat fishing industry on the Texas Gulf Coast,” by Robert B. Ditton, Richard N. Jarman, and Steve A. Woods, 8:1
Anaerobic bacteria
disease agents in fish, 10:10
“Anaerobic bacteria as possible disease agents in fish,” by Lanny R. Udey, 10:10
“Annotated bibliography on lobster trapping and related subjects,” by Ronald Joel Smolowitz, 5:6-68
Argopecten gibbus—see Scallop, calico
“(The) Atlantic halibut and its utilization,” by J. R. T. Nickerson, 7:21
Atwater, W. O.
chemical composition of fish, 7:1
Austen, Jack D.—see McConnell and Austen

B

- Bacterial diseases—see Diseases, bacterial
“Bacterial diseases of fishes and their control,” by S. F. Snieszko, 10:2
“Bacteriologic studies of Long Island shellfish hatcheries: An abstract,” by Louis Leibovitz, 10:8
Balaena mysticetus—see Whale, bowhead
Ballard, Kenneth, and Vito Blomo, “Estimating the structure of capacity utilization in the fishing industry,” 8:29
Barnett, Harold J., Richard W. Nelson, Patrick J. Hunter, and Herman Groninger, “Use of carbon dioxide dissolved in refrigerated brine for the preservation of pink shrimp (*Pandalus* spp.),” 9:24
Barrow, Alaska
whale, bowhead
1976 catch by Eskimos, 11:23, 25
Bass, striped
Pacific Northwest commercial fishery, 1922-74
commercial fishery, 1:18
fish introduction and range extension, 1:18
species description, 1:18
trends, 1:20

- Bauersfeld, Paul E., Jr.—see Hale and Bauersfeld
Bayer, Robert C., Margie Lee Gallagher, and Dale F. Leavitt, “Nutrient requirement of the lobster and nutrition pathology,” 10:44
Blomo, Vito—see Ballard and Blomo
Nichols, “Catch-effort and price-cost trends in the Gulf of Mexico shrimp fishery: Implications on Mexico's extended jurisdiction,” 8:24
Blott, Alan J., “A preliminary study of timed release mechanisms for lobster traps,” 5:6-44
Blue, Jack L.—see Gratzek et al.
“Blue crab larval culture: Methods and management,” by Mark R. Millikin, 11:10
Bowser, P. R.—see Plumb and Bowser
British Columbia
American shad, history of commercial fishery, 2:29
Brown, Albert L., “Immunity and practical vaccine development,” 3:2
Brown, C.—see Robohm and Brown
Busch, R. A., “Enteric red mouth disease (Hagerman Strain),” 3:42
Buzzell, David H.—see Sidwell et al.

C

- Caillouet, Charles W., and Frank J. Patella, “Relationship between size composition and ex-vessel value of reported shrimp catches from two Gulf Coast States with different harvesting strategies,” 2:14
California
oceanographic conditions, summer 1977, 2:24
Callinectes sapidus—see Crab, blue
Carver, J. H.—see Ronsivalli et al.
“Catch-effort and price-cost trends in the Gulf of Mexico shrimp fishery: Implications on Mexico's extended jurisdiction,” by Vito Blomo, Wade L. Griffin, and John P. Nichols, 8:24
Catfish, channel
immunization against two bacterial diseases, 3:18
serological screening of virus, 3:30
virus disease epizootiology
cell culture characteristics, 3:28
clinical signs and pathogenesis, 3:27
control, 3:29
immunological response, 3:28
morphology and biophysical nature, 3:28
transmission, 3:26
virus replication
phosphonoacetic acid inhibition of, 10:24
virus research, recent developments, 10:12
“Cellular immunity in fish as measured by lymphocyte stimulation,” by M. Michael Sigel, John C. Lee, E. Churchill McKinney, and Diana M. Lopez, 3:6
Celtic Sea
herring stocks
status, fished by the Federal Republic of Germany fleet, 4:22
Channel catfish virus disease—see Catfish, channel
“Characteristics, participation, and motivations of Texas charter boat fishermen,” by Robert B.

- Ditton, Thomas J. Mertens, and Martin P. Schwartz, 8:8
“Charles Woods, pioneer scientist on chemical composition of fish,” by Maurice E. Stansby, 7:1
Chelonia mydas—see Turtle, green sea
Cheng, Thomas C., “Larval nematodes parasitic in shellfish,” 10:39
Clifford, R. A.—see Pecci et al.
Clupea harengus—see Herring roe
Cod, Atlantic
 piscine erythrocytic necrosis (PEN), 10:16
“Comments on trends in research on parasitic diseases of shellfish and fish,” by Victor Sprague, 10:26
“Composition of the edible portion of raw (fresh or frozen) crustaceans, finfish, and mollusks. III. Microelements,” by Virginia D. Sidwell, Audrey L. Loomis, Karen J. Loomis, Pauline R. Foncannon, and David H. Buzzell, 9:1
“Composition of the edible portion of raw (fresh or frozen) crustaceans, finfish, and mollusks. IV. Vitamins,” by Virginia D. Sidwell, Audrey L. Loomis, Pauline R. Foncannon, and David H. Buzzell, 12:1
“(A) concept for assuring the quality of seafoods to consumers,” by L. J. Ronsivalli, C. Gorga, J. D. Kaylor, and J. H. Carver, 1:1
Contaminants—see Pollutants
“Control of fish diseases,” by S. F. Snieszko, 3:65
Cooper, R. A.—see Pecci et al.
Crab, blue
larval culture
egg development in vitro, 11:11
egg mass: age and disease, 11:11
feed organisms, culture, 11:13
feeding, 11:12
holding tanks for mass culture, 11:13
mating and spawning, 11:10
survival in natural vs artificial systems, 11:10
water quality, characteristics, and optimal parameters, 11:11
viral diseases of, 10:13
Crab, portunid
Portunus spinimanus
U.S. Virgin Islands, resource potential in, 7:12
Creamer, Donn L.—see Slabyj et al.
Crustaceans
composition of edible portion of raw (fresh or frozen)
aluminum, 9:1
arsenic, 9:1
ascorbic acid, 12:1
barium, 9:1
biotin, 12:1
cadmium, 9:1
choline, 12:1
chromium, 9:1
cobalt, 9:1
copper, 9:1
fluorine, 9:1
folic acid, 12:1
inositol, 12:1
iodine, 9:1

- iron, 9:1
lead, 9:1
manganese, 9:1
mercury, 9:1
mercury, organic, 9:1
molybdenum, 9:1
niacin, 12:1
nickel, 9:1
pantothenic acid, 12:1
pyridoxine, 12:1
riboflavin, 12:1
selenium, 9:1
silver, 9:1
thiamin, 12:1
tin, 9:1
vanadium, 9:1
vitamin A, 12:1
vitamin B₁₂, 12:1
vitamin D, 12:1
vitamin E, 12:1
zinc, 9:1
- Crustaceans, decapod—see Decapods
- D**
- Damann, Arthur E.—see Olsen et al.
- Decapods
 protozoan diseases of, 10:32
- Disease
 lobster
 Homarus americanus, 10:14
- Diseases
 environmentally related
 fish and shellfish, 10:43
 teleost liver, PCB-induced alterations in
 model for environmental disease in fish,
 10:47
- Diseases, bacterial
 anaerobic
 disease agents in fish, 10:10
- fish
 control of, 10:2
 shellfish hatcheries, Long Island
 bacteriologic studies, 10:8
- Diseases, parasitic
 shellfish
 larval nematodes in, 10:39
- shellfish and fish
 research, comments on trends, 10:26
- warm-water fishes, 10:36
- Diseases, protozoan
 crustaceans, decapod, 10:32
- "Diseases and defense mechanisms of the lobster, *Homarus americanus*," by James E. Stewart, 10:4
- Ditton, Robert B., Richard N. Jarman, and Steve A. Woods, "An analysis of the charter boat fishing industry on the Texas Gulf Coast," 8:1
- _____, Thomas J. Mertens, and Martin P. Schwartz, "Characteristics, participation, and motivations of Texas charter boat fishermen," 8:8
- Dornheim, Holger, "Status of the herring stocks fished by the Federal Republic of Germany fleet," 4:21
- E**
- "Economic aspects of the development of the herring roe gillnet fishery, southeastern Alaska, 1976," by Howard O. Ness, 4:25
- Eels
 catching and preparation methods, 4:1
 life history, 4:1
 world demand, 4:1
- "Eels and their utilization," by J. Perry Lane, 4:1
- "(The) effects of different escape vents on the selectivity of lobster traps," by Vernon E. Nulka, 5:6:50
- Elliott, Diane, "Fish disease inspection and certification," 3:69
- Enteric red mouth disease
 Hagerman strain
 agglutination test, passive, 3:47
 asymptomatic disease carrier state, 3:45
 carrier identification, 3:47
 carrier infections, cyclical nature, 3:47
 causative organism, 3:43
 clinical signs and pathology, 3:43
 economic effects, 3:43
 epidemiology, 3:42
 induced humoral immune response, 3:49
 mortality and transmission, 3:43
- "Enteric red mouth disease (Hagerman strain)," by R. A. Busch, 3:42
- "Environmental parameters affecting fish physiology in water reuse systems," by Thomas L. Meade, 10:46
- "Environmentally related diseases of marine fish and shellfish," by Carl J. Sindermann, 10:43
- "Epizootiology of channel catfish virus disease," by John A. Plumb, 3:26
- "Estimating the structure of capacity utilization in the fishing industry," by Kenneth Ballard and Vito Blomo, 8:29
- F**
- Farley, C. Austin, "Virus and viruslike lesions in marine mollusks," 10:18
- Federal Republic of Germany
 herring stocks, status, 4:21
- Finfish
 composition of edible portion of raw (fresh or frozen)
 aluminum, 9:1
 arsenic, 9:1
 ascorbic acid, 12:1
 barium, 9:1
 biotin, 12:1
 cadmium, 9:1
 choline, 12:1
 chromium, 9:1
 cobalt, 9:1
 copper, 9:1
 fluorine, 9:1
 folic acid, 12:1
 inositol, 12:1
 iodine, 9:1
 iron, 9:1
 lead, 9:1
 manganese, 9:1
 mercury, 9:1
 mercury, organic, 9:1
 molybdenum, 9:1
 niacin, 12:1
 nickel, 9:1
 pantothenic acid, 12:1
 pyridoxine, 12:1
 riboflavin, 12:1
 selenium, 9:1
 silver, 9:1
 thiamin, 12:1
 tin, 9:1
 vanadium, 9:1
 vitamin A, 12:1
 vitamin B₁₂, 12:1
 vitamin D, 12:1
 vitamin E, 12:1
 zinc, 9:1
- Fish
 bacterial disease, control of, 10:2
 chemical composition of, 7:1
 preparation, 1:5
- Fish, freshwater
 parasitic diseases
 cestodes, 3:57
 crustaceans, 3:57
 nematodes, 3:57
- protozoans, 3:56
 trematodes, 3:57
- Fish, ornamental
 diseases and problems
 bacteriological examination and results, 3:58, 59
 parasitological examination and results, 3:58, 59
 virological examination and results, 3:59, 60
- Fish, warmwater
 disease incidence on farms in south-central United States
 antibiotics used as prophylactics, 3:40
 bacterial infection and oxygen depletion incidence, 3:38
 disease incidence, 3:38
 protozoan infection incidence, 3:38
 treatment delivery system, 3:40
- parasitic diseases, 10:36
- "Fish disease inspection and certification," by Diane Elliott, 3:69
- Fish diseases
 anaerobic bacteria, possible disease agents, 10:10
 bacterial, control of, 10:2
- control
 chemoprophylaxis, 3:67
 chemotherapy effectiveness, 3:67
 drug administration, 3:65
 drugs commonly used, 3:65
 parasite control, 3:67
 relationship of host, pathogen, and environment, 3:65
- environmentally related, 10:43
- inspection and certification
 bacterial, 3:70
 diagnostic examination of fish, 3:71
 parasitic, 3:70
 sampling, 3:69
 viral, 3:69
- model for environmental
 teleost liver, PCB-induced alterations in, 10:47
- parasitic, 10:36
- parasitic
 research, comments on trends, 10:26
- Fish flavors
 petroleum pickup
 nature, 1:13
 research, 1:14
- Fish immunization
 Aeromonas salmonicida bacterins
 aggregating and nonaggregating strains, 3:12
 vaccines, 3:15
 virulence of strains, 3:13
- Fish physiology
 environmental parameters affecting
 water reuse systems, 10:46
- Fish traps
 West Indian, mesh selectivity of, 7:15
- Fishermen, Texas charter boat
 characteristics, participation, and motivations
 fishing motivations, analysis, 8:10
 fishing motivations, understanding, 8:10
 group types, 8:10
 motives, importance, 8:11
 population, study, 8:9
 recreational fishing backgrounds and participation, 8:9
 sociodemographic characteristics, 8:9
- Fishing industry
 estimating structure of capacity utilization
 economic capacity, 8:29
 engineering capacity, 8:29
 harvesting sector, 8:31
 price-capacity utilization relationships, 8:32

processing sector, 8:31
theoretical background, 8:30
Fishing industry, charter boat
Texas Gulf Coast
business characteristics, 8:3
financial information, 8:5
fishing activities, 8:4
operators, 8:3
organization, 8:2
Flatfish
ulcer cause, presumptive *Vibrio* species, 10:5
"Flavors in fish from petroleum pickup," by
Maurice E. Stansby, 1:13
Foncannon, Pauline R.—see Sidwell et al.
Fryer, J. L.—see Udey and Fryer
—, J. S. Rohovec, and R. L. Garrison,
"Immunization of salmonids for control of
vibriosis," 3:20
Furunculosis
Puget Sound, Washington
salmon, marine cultured, 3:52

G

Gallagher, Margie Lee—see Bayer et al.
Gardner, George R., "A review of histopatholog-
ical effects of selected contaminants on some
marine organisms," 10:51
Garrison, R. L.—see Fryer et al.
Georges Bank
herring stocks
status, fished by the Federal Republic of
Germany fleet, 4:23
sea scallop, resources 1975
distribution and abundance, 2:20
length frequency, 2:22
spawning season, 2:20
"Ghost fishing of vented and unvented lobster,
Homarus americanus, traps," by K. J. Pecci,
R. A. Cooper, C. D. Newell, R. A. Clifford,
and R. J. Smolowitz, 5:6-9
Gorga, L. J.—see Ronsivalli et al.
Gratzek, John B., Emmett B. Shotts, and Jack L.
Blue, "Ornamental fish: Diseases and prob-
lems," 3:58
Griffin, Wade L.—see Blomo et al.
Groninger, Herman—see Barnett et al.
Gulf Coast
shrimp catches, Texas and Louisiana
size composition, effect on ex-vessel value,
2:14
Gulf of Mexico
shrimp fishery, catch-effort and price-cost
trends
Mexico, implications on extended jurisdic-
tion, 8:24

H

Haines, Harold, "A herpesvirus disease of green
sea turtles in aquaculture," 3:33
Hale, Malcolm B., and Paul E. Bauersfeld, Jr.,
"Preparation of a menhaden hydrolysate for
possible use in a milk replacer," 8:14
Halibut, Atlantic
utilization, 7:21
Harrell, Lee W., "Vibriosis and current salmon
vaccination procedures in Puget Sound, Wash-
ington," 3:24
Hebrides
herring stocks
status, fished by the Federal Republic of
Germany fleet, 4:23
"(A) herpesvirus disease of green sea turtles in
aquaculture," by Harold Haines, 3:33
Herring
status, fished by Federal Republic of Germany
fleet, 4:21
Herring roe
gillnet fishery, southeastern Alaska, 1976
economic aspects of development, 4:25
Hinton, David E., James E. Klaunig, and

Michael M. Lipsky, "PCB-induced alterations
in teleost liver: A model for environmental
disease in fish," 10:47
Hippoglossus hippoglossus—see Halibut, Atlan-
tic
Homarus americanus—see Lobster, American
Hunter, Patrick J.—see Barnett et al.
Hutchinson, Roger W., "Status and problems of
the American shrimp industry," 9:29

I

Ictalurus punctatus—see Catfish, channel
Illex illecebrosus—see Squid, shortfin
Immunity
vaccine development, 3:2
Immunity, cellular
fish, as measured by lymphocyte stimulation
evidence that antibodies can regulate lym-
phocyte responses, 3:8
immune responses, regulation, 3:7
immunoglobulins, 3:7
immunologic response, 3:7
suppressor cells, 3:10
"Immunity and practical vaccine development,"
by Albert L. Brown, 3:2
"Immunization of channel catfish, *Ictalurus
punctatus*, against two bacterial diseases," by
John H. Schachte, Jr., 3:18
"Immunization of fish with bacterins of *Aero-
monas salmonicida*," by L. R. Udey and J. L.
Fryer, 3:12
"Immunization of salmonids for control of vib-
riosis," by J. L. Fryer, J. S. Rohovec, and R.
L. Garrison, 3:20
"Incidence of disease in warmwater fish farms in
the south-central United States," by F. P.
Meyer, 3:38
Ingraham, W. James, Jr., and Cuthbert M. Love,
"Oceanographic conditions off California to
Vancouver Island in the summer of 1977,"
2:24

J

Jarman, Richard N.—see Ditton et al.
Johnson, Phyllis T., "Viral diseases of the blue
crab, *Callinectes sapidus*," 10:13

K

Kaktovik, Alaska
whale, bowhead
1976 catch by Eskimos, 11:27
Kaylor, J. D.—see Ronsivalli et al.
Kivalina, Alaska
whale, bowhead
1976 catch by Eskimos, 11:22
Klaunig, James E.—see Hinton et al.
Koment, Roger W., "Phosphonoacetic acid in-
hibition of channel catfish virus replication,"
10:24
Lane, J. Perry, "Eels and their utilization," 4:1
LaPlace, Joseph A.—see Olsen et al.
"Larval nematodes parasitic in shellfish," by
Thomas C. Cheng, 10:39
Leavitt, Dale F.—see Bayer et al.
Lee, John C.—see Sigel et al.
Leibovitz, Louis, "Bacteriologic studies of Long
Island shellfish hatcheries: An abstract," 10:8
—, "Shellfish diseases," 3:61
Lind, H. F.—see Lux et al.
Lipsky, Michael M.—see Hinton et al.
Lobster
Homarus americanus
diseases and defense mechanisms of, 10:4
nutrient requirement and nutrition pathology,
10:44
Lobster, American
ghost fishing of vented and unvented traps,
5:6-9

L

Long Island
shellfish hatcheries
bacteriologic studies, 10:8
Loomis, Audrey L.—see Sidwell et al.
Loomis, Karen J.—see Sidwell et al.
Lopez, Diana M.—see Sigel et al.
Louisiana
shrimp catches
size composition, effect on ex-vessel value,
2:14
Love, Cuthbert M.—see Ingraham and Love
Lux, F. E., J. R. Uzman, and H. F. Lind,
"Strandings of shortfin squid, *Illex illece-
brosus*, in New England in fall 1976," 1:21

M

McAllister, Philip E., "Viruses and virus dis-
eases of salmonid fishes," 10:21
McConnell, Stewart, and Jack D. Austen,
"Serological screening of channel catfish
virus," 3:30
MacLean, Sharon A.—see Sawyer and MacLean
MacKenzie, Clyde L., Jr., Arthur S. Merrill, and
Fredric M. Serchuk, "Sea scallop resources
off the northeastern U.S. coast," 2:19
McKinney, E. Churchill—see Sigel et al.
Maine
salmonids
vibriosis in, 10:9
Marquette, Willman M., "The 1976 catch of
bowhead whales, *Balaena mysticetus*, by
Alaskan Eskimos," 11:18
Meade, Thomas L., "Environmental parameters
affecting fish physiology in water reuse sys-
tems," 10:46
Menhaden
hydrolysate
preparation of, for possible use in a milk
replacer, 8:14
Merrill, Arthur S.—see MacKenzie et al.
Mertens, Thomas J.—see Ditton et al.
"Mesh selectivity of West Indian fish traps," by
David A. Olsen, Arthur E. Dammann, and
Joseph A. LaPlace, 7:15
Mexico
catch-effort and price-cost trends in Gulf of
Mexico shrimp fishery
extended jurisdiction implications, 8:24
Meyer, F. P., "Incidence of disease in warmwa-
ter fish farms in the south-central United
States," 3:38
Middle Atlantic Shelf
sea scallop, resources 1975
distribution and abundance, 2:21
length frequency, 2:22
Milk replacer
preparation of menhaden hydrolysate for pos-
sible use in, 8:14
Milliken, Mark R., "Blue crab larval culture:
Methods and management," 11:10
Mollusks
composition of edible portion of raw (fresh or
frozen)
aluminum, 9:1
arsenic, 9:1
ascorbic acid, 12:1
barium, 9:1
biotin, 12:1
cadmium, 9:1
choline, 12:1
chromium, 9:1
cobalt, 9:1
copper, 9:1
fluorine, 9:1
folic acid, 12:1
inositol, 12:1
iodine, 9:1
iron, 9:1
lead, 9:1
manganese, 9:1

mercury, 9:1
 mercury, organic, 9:1
 molybdenum, 9:1
 niacin, 12:1
 nickel, 9:1
 pantothenic acid, 12:1
 pyridoxine, 12:1
 riboflavin, 12:1
 selenium, 9:1
 silver, 9:1
 thiamin, 12:1
 tin, 9:1
 vanadium, 9:1
 vitamin A, 12:1
 vitamin B₁₂, 12:1
 vitamin D, 12:1
 vitamin E, 12:1
 zinc, 9:1

Mollusks, marine
 viruses and viruslike lesions, 10:18
Morone saxatilis—see Bass, striped
 Mussel, blue
 cultivated and natural stocks
 seasonal effect on yield, proximate composition, and quality of meats, 8:18
Mytilus edulis—see Mussel, blue

N

Nelson, Richard W.—see Barnett et al.
 Nematodes
 larval, parasitic in shellfish, 10:39
 Ness, Howard O., "Economic aspects of the development of the herring roe gillnet fishery, southeastern Alaska, 1976," 4:25
 "(A) new bacterium (presumptive *Vibrio* species) causing ulcers in flatfish," by R. A. Robohm and C. Brown, 10:5
 New Hampshire
 salmonids
 vibriosis in, 10:9
 Newell, C. D.—see Pecci et al.
 Nichols, John P.—see Blomo et al.
 Nicholson, Bruce L.—see Reno et al.
 Nickerson, J. T. R., "The Atlantic halibut and its utilization," 7:21
 "(The) 1976 catch of bowhead whales, *Balaena mysticetus*, by Alaskan Eskimos," by Willman M. Carquette, 11:18
 North Sea
 herring stocks
 status, fished by the Federal Republic of Germany fleet, 4:21
 Norwegian Sea
 herring stocks
 status, fished by the Federal Republic of Germany fleet, 4:23
 Novotny, Anthony J., "Vibriosis and furunculosis in marine cultured salmon in Puget Sound, Washington," 3:52
 Nuiqsut, Alaska
 whale, bowhead
 1976 catch by Eskimos, 11:27
 Nulk, Vernon E., "The effects of different escape vents on the selectivity of lobster traps," 5:6:50
 "Nutrient requirement of the lobster and nutrition pathology," by Robert C. Bayer, Margie Lee Gallagher, and Dale F. Leavitt, 10:44
 Nutrition
 lobster
 requirement and pathology, 10:44

O

"Oceanographic conditions off California to Vancouver Island in the summer of 1977," by W. James Ingraham, Jr. and Cuthbert M. Love, 2:24
 O'Gradnick, Joseph, "Susceptibility studies of various salmonids to whirling disease: Histological staining and spore concentration pro-

cedures," 10:30
 Olsen, David A., Arthur E. Dammann, and Joseph A. LaPlace, "Mesh selectivity of West Indian fish traps," 7:15
 ———, ———, and ———, "Portunus spinimanus Latreille, a portunid crab with resource potential in the U.S. Virgin Islands," 7:12
 Oregon
 American shad, history of commercial fishery, 2:29
 oceanographic conditions, summer 1977, 2:24
 "Ornamental fish: Diseases and problems," by John B. Gratzek, Emmett B. Shotts, and Jack L. Blue, 3:58
 Overstreet, Robin M., "Trypanorhynch infections in the flesh of sciaenid fishes," 10:37

P

Pacific Northwest
 American shad, history of commercial fishery, 2:29
 sturgeon, commercial fishery, 7:17
 "(The) Pacific Northwest commercial fishery for American shad," by Norman B. Parks, 2:29
 "(The) Pacific Northwest commercial fishery for striped bass, 1922-74," by Norman B. Parks, 1:18
 "(The) Pacific Northwest commercial fishery for sturgeon," by Norman B. Parks, 7:17
 "Parasitic diseases of freshwater fishes," by W. A. Rogers, 3:56
 Parasitic diseases—see Diseases, parasitic
 Parks, Norman B., "The Pacific Northwest commercial fishery for American shad," 2:29
 ———, "The Pacific Northwest commercial fishery for striped bass, 1922-74," 1:18
 ———, "The Pacific Northwest commercial fishery for sturgeon," 7:17
 Patella, Frank J.—see Caillouet and Patella
Patinopecten caurinus—see Scallop, Pacific sea
 "PCB-induced alterations in teleost liver: A model for environmental disease in fish," by David E. Hinton, James E. Klaunig, and Michael M. Lipsky, 10:47
 Pecci, K. J., R. A. Cooper, C. D. Newell, R. A. Clifford, and R. J. Smolowitz, "Ghost fishing of vented and unvented lobster, *Homarus americanus*, traps," 5:6:9
Penaeus aztecus—see Shrimp, brown
Penaeus setiferus—see Shrimp, white
 Peters, John A., "Scallops and their utilization," 11:1
 Philippon-Fried, Marie—see Reno et al.
 Phosphonoacetic acid
 inhibition of virus replication in channel catfish, 10:24
 "Phosphonoacetic acid inhibition of channel catfish virus replication," by Roger W. Koment, 10:24
 Piscine erythrocytic necrosis
 cod, Atlantic, 10:16
 "Piscine erythrocytic necrosis (PEN): A viral infection of the Atlantic cod and other marine fishes," by Paul Reno, Marie Philippon-Fried, Bruce L. Nicholson, and Stuart Sherburne, 10:16
Placopecten magellanicus—see Scallop, sea
 Plumb, John A., "Epizootiology of channel catfish virus disease," 3:26
 ———, and P. R. Bowser, "Recent developments in channel catfish virus research," 10:12
 Point Hope, Alaska
 whale, bowhead
 1976 catch by Eskimos, 11:22
 Pollutants
 histopathological effects on marine organisms, 10:51
 "Portunus spinimanus Latreille, a portunid crab

with resource potential in the U.S. Virgin Islands," by David A. Olsen, Arthur E. Dammann, and Joseph A. LaPlace, 7:12
Portunus spinimanus—see Crab, portunid
 "(A) preliminary study of timed release mechanisms for lobster traps," by Alan J. Blott, 5:6:44
 "Preparation of a menhaden hydrolysate for possible use in a milk replacer," by Malcolm B. Hale and Paul E. Bauersfeld, Jr., 8:14
 "Preparation of fish fillet blocks," by John J. Ryan, 1:5
 "(The) principal parasitic diseases of warm-water fishes," by W. A. Rogers, 10:36
 Protozoan diseases—see Diseases, protozoan
 Puget Sound, Washington
 vibriosis and current salmon vaccination procedures, 3:24

R

"Recent developments in channel catfish virus research," by J. A. Plumb and P. R. Bowser, 10:12
 "Relationship between size composition and ex-vessel value of reported shrimp catches from two Gulf Coast States with different harvesting strategies," by Charles W. Caillouet and Frank J. Patella, 2:14
 Reno, Paul, Marie Philippon-Fried, Bruce L. Nicholson, and Stuart Sherburne, "Piscine erythrocytic necrosis (PEN): A viral infection of the Atlantic cod and other marine fishes," 10:16
 "(A) review of histopathological effects of selected contaminants on some marine organisms," by George R. Gardner, 10:51
 "Review of oxygen depletion and associated mass mortalities of shellfish in the Middle Atlantic Bight in 1976," by Frank W. Steimle and Carl J. Sindermann, 12:17
 Robohm, R. A., and C. Brown, "A new bacterium (presumptive *Vibrio* species) causing ulcers in flatfish," 10:5
 Rockland, Steven, "The San Diego tuna industry and its employment impact on the local economy," 7:5
 Rogers, W. A., "Parasitic diseases of freshwater fishes," 3:56
 ———, "The principal parasitic diseases of warm-water fishes," 10:36
 Rohovec, J. S.—see Fryer et al.
 Ronsivalli, Louis J., "Sharks and their utilization," 2:1
 ———, C. Gorga, J. D. Kaylor, and J. H. Carver, "A concept for assuring the quality of seafoods to consumers," 1:1
 Ryan, John J., "Preparation of fish fillet blocks," 1:5

S

St. Lawrence Island, Alaska
 whale, bowhead
 1976 catch by Eskimos, 11:21
 Salmon, marine cultured
 vibriosis and furunculosis, Puget Sound, Washington
 description of diseases, 3:52
 diagnostic procedures, 3:53
 disease observations, 1972-73, 3:53
 drug sensitivity tests, 3:53
 mortality assessments, 3:53
 therapeutic treatment procedures, 3:53
 Salmonids
 immunization for control of vibriosis
 bacterin, length of oral administration, 3:21, 22
 bacterin administration and challenge, 3:20
 bacterin concentration, 3:21, 22
 bacterin preparation, 3:20
 oral and injected bacterins, comparison,

- 3:21, 22
temperature effect on oral immunization, 3:21, 22
- Puget Sound, Washington
vibriosis and vaccination procedures, 3:24
viruses and virus diseases of, 10:21
whirling disease, susceptibility studies, 10:30
- San Diego
tuna industry
employment impact on the local economy, 7:5
“(The) San Diego tuna industry and its employment impact on the local economy,” by Steven Rockland, 7:5
- Sawyer, Evelyn S., “Vibriosis in Maine and New Hampshire salmonids,” 10:9
- Sawyer, Thomas K., and Sharon A. MacLean, “Some protozoan diseases of decapod crustaceans,” 10:32
- Scallop, bay
utilization
aquaculture, 11:9
biology, 11:3
harvesting, 11:5
history, 11:1
processing, 11:6, 7
- Scallop, calico
utilization
aquaculture, 11:9
biology, 11:3
harvesting, 11:5
history, 11:2
processing, 11:6, 7
- Scallop, Pacific sea
utilization
biology, 11:4
history, 11:2
- Scallop, sea
resources, northeastern U.S. coast 1975, 2:19
utilization
aquaculture, 11:8
biology, 11:2
harvesting, 11:4
history, 11:1
processing, 11:6
quality problems, 11:8
“Scallops and their utilization,” by John A. Peters, 11:1
- Schachte, John H., Jr., “Immunization of channel catfish, *Ictalurus punctatus*, against two bacterial diseases,” 3:18
- Schwartz, Martin P.—see Ditton et al.
- Sciaenids
trypanorhynch infections in, 10:37
“Sea scallop resources off the northeastern U.S. coast,” by Clyde L. MacKenzie, Jr., Arthur S. Merrill, and Fredric M. Serchuk, 2:19
- Seafood
concept for assuring quality to consumers, 1:1
“Seasonal effect on yield, proximate composition, and quality of blue mussel, *Mytilus edulis*, meats obtained from cultivated and natural stock,” by Bohdan M. Slabyj, Donn L. Creamer, and Ruth H. True, 8:18
- Seidel, Wilber R., and John W. Watson, Jr., “A trawl design: Employing electricity to selectively capture shrimp,” 9:21
- Serchuk, Fredric M.—see MacKenzie et al.
“Serological screening of channel catfish virus,” by Stewart McConnell and Jack D. Austen, 3:30
- Shad, American
Pacific Northwest commercial fishery
history, 2:29
- Sharks
description and physical characteristics, 2:1
occurrence and habits, 2:1
relationship to man, 2:1
utilization, 2:1
“Sharks and their utilization,” by Louis J. Ron-sivalli, 2:1
- Shellfish
oxygen depletion and mass mortalities in Middle Atlantic Bight in 1976
fish and shellfish stocks, 12:19
fish populations, effects on, 12:22
meteorological and hydrological events, 12:19
oxygen depletion, extent, 12:18
shellfish, effects on, 2:20
- Shellfish diseases
arthropods, 3:62
bacterial, 3:62
environmentally related, 10:43
etiology unknown, 3:62
fungal, 3:62
hatchery operation studies, 3:62
helminthic, 3:62
larval nematodes parasitic in, 10:39
parasitic, 3:62, 10:26
testing bacterial pathogenicity, 3:63
viral, 3:61
“Shellfish diseases,” by Louis Leibovitz, 3:61
- Shellfish hatcheries
Long Island
bacteriologic studies, 10:8
Sherburne, Stuart—see Reno et al.
Shotts, Emmett B.—see Gratzek et al.
- Shrimp
catches, two Gulf States
relationship between size composition and ex-vessel value, 2:14
- Shrimp, brown
Texas and Louisiana catches
size composition, effect on ex-vessel value, 2:14
- Shrimp, pink
preserved in carbon dioxide dissolved in refrigerated brine
bacteriological measurements, 9:25
CO₂ concentration, 9:27
pH, 9:26
refrigerated brine equipment, 9:25
salt concentration, 9:27
sensory evaluation, 9:27
- Shrimp, white
Texas and Louisiana catches
size composition, effect on ex-vessel value, 2:14
- Shrimp fishery
catch-effort and price-cost trends in Gulf of Mexico
implications on Mexico's extended jurisdiction, 8:24
- Shrimp industry, American
status and problems
jurisdictions, extended, 9:31
imports, 9:30
insurance rates, 9:31
operating costs, 9:30
overcapitalization, 9:30
- Sidwell, Virginia D., Audrey L. Loomis, Karen J. Loomis, Pauline R. Foncannon, and David H. Buzzell, “Composition of the edible portion of raw (fresh or frozen) crustaceans, finfish, and mollusks. III. Microelements,” 9:1
_____, Pauline R. Foncannon, and David H. Buzzell, “Composition of the edible portion of raw (fresh or frozen) crustaceans, finfish, and mollusks. IV. Vitamins,” 12:1
- Sigel, M. Michael, John C. Lee, E. Churchill McKinney, and Diana M. Lopez, “Cellular immunity in fish as measured by lymphocyte stimulation,” 3:6
- Sindermann, Carl J., “Environmentally related diseases of marine fish and shellfish,” 10:43
—see also Steimle and Sindermann
- Slabyj, Bohdan M., Donn L. Creamer, and Ruth H. True, “Seasonal effect on yield, proximate composition, and quality of blue mussel, *Mytilus edulis*, meats obtained from cultivated and natural stock,” 8:18
- Smolowitz, Ronald Joel, “Annotated bibliography on lobster trapping and related subjects,” 5-6:68
_____, “Trap design and ghost fishing: An overview,” 5-6:2
_____, “Trap design and ghost fishing: Discussion,” 5-6:59
—see also Pecci et al.
- Snieszko, S. F., “Bacterial diseases of fishes and their control,” 10:2
_____, “Control of fish diseases,” 3:65
“Some protozoan diseases of decapod crustaceans,” by Thomas K. Sawyer and Sharon A. MacLean, 10:32
- Sprague, Victor, “Comments on trends in research on parasitic diseases of shellfish and fish,” 10:26
- Squid, shortfin
strandings in New England in fall 1976
catches by research vessel, 1:22
distribution and fishery, 1:21
estimating numbers stranded, 1:25
examining stranded squid, 1:23
reasons for strandings, 1:25
stranding reports, 1:22
strandings, past, 1:23
- Stansby, Maurice E., “Charles Woods, pioneer scientist on chemical composition of fish,” 7:1
_____, “Flavors in fish from petroleum pickup,” 1:13
“Status and problems of the American shrimp industry,” by Roger W. Hutchinson, 9:29
“Status of the herring stocks fished by the Federal Republic of Germany fleet,” by Holger Dornheim, 4:21
- Steimle, Frank W., and Carl J. Sindermann, “Review of oxygen depletion and associated mass mortalities of shellfish in the Middle Atlantic Bight in 1976,” 12:17
- Stewart, James E., “Diseases and defense mechanisms of the lobster, *Homarus americanus*,” 10:4
“Strandings of shortfin squid, *Illex illecebrosus*, in New England in fall 1976,” by F. E. Lux, J. R. Uzmman, and H. F. Lind, 1:21
- Sturgeon
Pacific Northwest commercial fishery, 7:17
“Susceptibility studies of various salmonids to whirling disease: Histological staining and spore concentration procedures,” by Joseph O'Grodnick, 10:30

T

- Teleosts
liver, PCB-induced alterations in, 10:47
- Texas
fishermen, charter boat
characteristics, participation, and motivation, 8:8
Gulf Coast
charter boat fishing industry, 8:1
shrimp catches
size composition, effect on ex-vessel value, 2:14
“Trap design and ghost fishing: An overview,” by Ronald Joel Smolowitz, 5-6:2
“Trap design and ghost fishing: Discussion,” by Ronald Joel Smolowitz, 5-6:59
- Traps, lobster
annotated bibliography on, and related sub-

jects, 5-6:68
 design and ghost fishing
 availability and catchability, 5-6:62
 bait, 5-6:63
 behavior, 5-6:63
 biological deterioration, 5-6:61
 catch escape panels, 5-6:65
 corrosion, 5-6:61
 destruction by dragged gear, 5-6:62
 effort, reduced, 5-6:65
 escape vents, 5-6:7
 escape vents, sublegal, 5-6:65
 escapement, 5-6:62
 factors affecting ghost fishing, 5-6:5
 fishery, 5-6:2
 habipots, 5-6:66
 head design, 5-6:60
 injury and mortality, 5-6:63
 lobster range, 5-6:2
 loss assesemnt, 5-6:64
 loss cutting, 5-6:65
 management, 5-6:3
 materials, 5-6:60
 pots, number lost, 5-6:59
 preventing and decreasing ghost fishing, 5-6:6
 sections, degradable, 5-6:65
 set-over days, 5-6:62
 shape, 5-6:59
 size, 5-6:59
 storm surge, 5-6:61
 substrate burial, 5-6:62
 escape vents
 ghost fishing, 5-6:11
 selectivity, 5-6:50
 ghost fishing of vented and unvented
 catch gear distribution, 5-6:10
 description of trawls and pots, 5-6:11
 environmental monitoring, 5-6:13
 ghost-pot catch, 5-6:17
 ghost-pot trawls, surveying, 5-6:12
 natural population, 5-6:13
 pot trawls, setting and hauling of surface-buoyed, 5-6:11
 study area, 5-6:9
 surface-hauled pot catch, 5-6:14
 timed release mechanisms
 release mechanism, 5-6:44, 45, 49
 release method, 5-6:44, 48
 release timer, 5-6:44, 45
 "(A) trawl design: Employing electricity to selectively capture shrimp," by Wilber R. Seidel and John W. Watson, Jr., 9:21
 Trawl, shrimp
 employing electricity to selectively capture electrical fishing technology, recent advances, 9:22
 webbing panel type separator trawls, 9:21
 True, Ruth H.—see Slabyj et al.
 Trypanorhynch infections
 sciaenid flesh, 10:37
 "Trypanorhynch infections in the flesh of sciaenid fishes," by Robin M. Overstreet, 10:37
 Tuna industry
 San Diego
 employment impact on the local economy, 7:5
 Turtle, green sea
 herpesvirus disease
 clinical and histological picture, 3:34
 farm facilities and operations, 3:33
 gray-patch disease development, 3:36
 isolation attempts, 3:37
 presence of virus in lesions, 3:35
 temperature effect, 3:36
 transmission studies, 3:35
 vaccination attempts with virus, 3:37
 virus origin, 3:36

U

Udey, Lanny R., "Anaerobic bacteria as possible disease agents in fish," 10:10
 —, and J. L. Fryer, "Immunization of fish with bacterins of *Aeromonas salmonicida*," 3:12
 United States
 northeastern coast, 1975
 sea scallop resources, 2:19
 west coast
 oceanographic conditions, summer 1977, 2:24
 United States, south-central
 warmwater fish farms
 disease incidence, 3:38
 "Use of carbon dioxide dissolved in refrigerated brine for the preservation of pink shrimp (*Penaeus* spp.)," by Harold J. Barnett, Richard W. Nelson, Patrick J. Hunter, and Herman Groninger, 9:24
 Uzmans, J. R.—see Lux et al.

V

Vaccine
 development and immunity
 bacterial, inactivated, 3:23
 bacterial, modified live, 3:3
 cell line, stable, 3:4
 fish vaccines, 3:4
 immunity type, 3:2
 inactivated vs. live, 3:2
 vaccine types, 3:2
 viral, 3:3
 Vancouver Island
 oceanographic conditions, summer 1977, 2:24
Vibrio species, presumptive
 ulcer cause in flatfish, 10:5
 Vibriosis
 immunization of salmonids for control, 3:20
 Puget Sound, Washington
 salmon, marine cultured, 3:52
 salmon vaccination procedures, 3:24
 salmonids
 Maine and New Hampshire, 10:9
 "Vibriosis and current salmon vaccination procedures in Puget Sound, Washington," by Lee W. Harrel, 3:24
 "Vibriosis and furunculosis in marine cultured salmon in Puget Sound, Washington," by Anthony J. Novotny, 3:52
 "Vibriosis in Maine and New Hampshire salmonids," by Evelyn S. Sawyer, 10:9
 Viral diseases
 catfish, channel
 phosphonoacetic acid inhibition of, 10:24
 Cod, Atlantic
 piscine erythrocytic necrosis (PEN), 10:16
 crab, blue, 10:13
 mollusks, marine, 10:18
 salmonids, 10:21
 "Viral diseases of the blue crab, *Callinectes sapidus*," by Phyllis T. Johnson, 10:13
 Virgin Islands
 Portunus spinimanus
 resource potential, 7:12
 Virus research
 channel catfish
 recent developments, 10:12
 "Viruses and virus diseases of salmonid fishes," by Philip E. McAllister, 10:21
 "Viruses and viruslike lesions in marine mollusks," by C. Austin Farley, 10:18

W

Wainwright, Alaska
 whale, bowhead
 1976 catch by Eskimos, 11:23

Wales, Alaska

 whale, bowhead
 1976 catch by Eskimos, 11:22
 Washington
 American shad, history of commercial fishery, 2:29
 oceanographic conditions, summer 1977, 2:24
 Water reuse systems
 environmental parameters affecting fish physiology, 10:46
 Watson, John W., Jr.—see Seidel and Watson
 West Indies
 fish traps, mesh selectivity of, 7:15
 Whale, bowhead
 1976 catch by Alaskan Eskimos
 autumn whaling, 11:25
 effort, 11:23
 spring whaling, 11:21
 utilization, 11:20
 whaling methods, 11:19
 Whirling disease
 salmonids, susceptibility studies of, 10:30
 Woods, Charles
 chemical composition of fish, 7:1
 Woods, Steve A.—see Ditton et al.

X, Y, Z

Xiphiidae—Swordfishes, 9:17

Papers in Marine Fisheries Review

Volume 40, Numbers 1-12, 1978

JANUARY

1276. "A Concept for Assuring the Quality of Seafoods to Consumers," L. J. Ronsivalli, C. Gorga, J. D. Kaylor, and J. H. Carver, 1:1-4.
1277. "Preparation of Fish Fillet Blocks," John J. Ryan, 1:5-12.
1278. "Flavors in Fish From Petroleum Pickup," Maurice E. Stansby, 1:13-17.
1279. "The Pacific Northwest Commercial Fishery for Striped Bass, 1922-74," Norman B. Parks, 1:18-20.
1280. "Strandings of Shortfin Squid, *Illex illecebrosus*, in New England in Fall 1976," F. E. Lux, J. R. Uzmans, and H. F. Lind, 1:21-26.

FEBRUARY

1281. "Sharks and Their Utilization," Louis J. Ronsivalli, 2:1-13.
1282. "Relationship Between Size Composition and Ex-Vessel Value of Reported Shrimp Catches From Two Gulf Coast States With Different Harvesting Strategies," Charles W. Caillouet and Frank J. Patella, 2:14-18.
1283. "Sea Scallop Resources off the Northeastern U.S. Coast, 1975," Clyde L. MacKenzie, Jr., Arthur S. Merrill, and Fredric M. Serchuk, 2:19-23.
1284. "Oceanographic Conditions off California to Vancouver Island in the Summer of 1977," W. James Ingraham, Jr. and Cuthbert M. Love, 2:24-28.

1285. "The Pacific Northwest Commercial Fishery for American Shad," Norman B. Parks, 2:29-31.

MARCH

1286. "Immunity and Practical Vaccine Development," Albert L. Brown, 3:2-5.
 1287. "Cellular Immunity in Fish as Measured by Lymphocyte Stimulation," M. Michael Sigel, John C. Lee, E. Churchill McKinney, and Diana M. Lopez, 3:6-11.
 1288. "Immunization of Fish With Bacterins of *Aeromonas salmonicida*," L. R. Udey and J. L. Fryer, 3:12-17.
 1289. "Immunization of Channel Catfish, *Ictalurus punctatus*, Against Two Bacterial Diseases," John H. Schachte, Jr., 3:18-19.
 1290. "Immunization of Salmonids for Control of Vibriosis," J. L. Fryer, J. S. Rohovec, and R. L. Garrison, 3:20-23.
 1291. "Vibriosis and Current Salmon Vaccination Procedures in Puget Sound, Washington," Lee W. Harrell, 3:24-25.
 1292. "Epizootiology of Channel Catfish Virus Disease," John A. Plumb, 3:26-29.
 1293. "Serological Screening of Channel Catfish Virus," Stewart McConnell and Jack D. Austen, 3:30-32.
 1294. "A Herpesvirus Disease of Green Sea Turtles in Aquaculture," Harold Haines, 3:33-37.
 1295. "Incidence of Disease in Warmwater Fish Farms in the South-Central United States," F. P. Meyer, 3:38-41.
 1296. "Enteric Red Mouth Disease (Hagerman Strain)," R. A. Busch, 3:42-51.
 1297. "Vibriosis and Furunculosis in Marine Cultured Salmon in Puget Sound, Washington," Anthony J. Novotny, 3:52-55.
 1298. "Parasitic Diseases of Freshwater Fishes," W. A. Rogers, 3:56-57.
 1299. "Ornamental Fish: Diseases and Problems," John B. Gratzek, Emmett B. Shotts, and Jack L. Blue, 3:58-60.
 1300. "Shellfish Diseases," Louis Leibovitz, 3:61-64.
 1301. "Control of Fish Diseases," S. F. Snieszko, 3:65-68.
 1302. "Fish Disease Inspection and Certification," Diane Elliott, 3:69-71.

APRIL

1303. "Eels and Their Utilization," J. Perry Lane, 4:1-20.
 1304. "Status of the Herring Stocks Fished by the Federal Republic of Germany Fleet," Holger Dornheim, 4:21-24.
 1305. "Economic Aspects of the Development of the Herring Roe Gillnet Fishery, Southeastern Alaska, 1976," Howard O. Ness, 4:25-30.

MAY-JUNE

1306. "Trap Design and Ghost Fishing: An Overview," Ronald Joel Smolowitz, 5:6-28.
 1307. "Ghost Fishing of Vented and Unvented Lobster, *Homarus americanus*, Traps," K. J. Pecci, R. A. Cooper, C. D. Newell, R. A. Clifford, and R. J. Smolowitz, 5:6-9-43.
 1308. "A Preliminary Study of Timed Release Mechanisms for Lobster Traps," Alan J. Blott, 5:6-44-49.
 1309. "The Effects of Different Escape Vents on the Selectivity of Lobster Traps," Vernon E. Nulak, 5:6-50-58.
 1310. "Trap Design and Ghost Fishing: Discussion," Ronald Joel Smolowitz, 5:6-59-67.
 1311. "Annotated Bibliography on Lobster Trapping and Related Subjects," Ronald Joel Smolowitz, 5:6-68-77.

JULY

1312. "Charles Woods, Pioneer Scientist on Chemical Composition of Fish," Maurice E. Stansby, 7:1-4.
 1313. "The San Diego Tuna Industry and Its Employment Impact on the Local Economy," Steven Rockland, 7:5-11.
 1314. "*Portunus spinimanus* Latreille, a Portunid Crab With Resource Potential in the U.S. Virgin Islands," David A. Olsen, Arthur E. Dammann, and Joseph A. LaPlace, 7:12-14.
 1315. "Mesh Selectivity of West Indian Fish Traps," David A. Olsen, Arthur E. Dammann, and Joseph A. LaPlace, 7:15-16.
 1316. "The Pacific Northwest Commercial Fishery for Sturgeon," Norman B. Parks, 7:17-20.
 1317. "The Atlantic Halibut and Its Utilization," J. T. R. Nickerson, 7:21-25.

AUGUST

1318. "An Analysis of the Charter Boat Fishing Industry on the Texas Gulf Coast," Robert B. Ditton, Richard N. Jarman, and Steve A. Woods, 8:1-7.
 1319. "Characteristics, Participation, and Motivations of Texas Charter Boat Fishermen," Robert B. Ditton, Thomas J. Mertens, and Martin P. Schwartz, 8:8-13.
 1320. "Preparation of a Menhaden Hydrolysate for Possible Use in a Milk Replacer," Malcolm B. Hale and Paul E. Bauersfeld, Jr., 8:14-17.
 1321. "Seasonal Effect on Yield, Proximate Composition, and Quality of Blue Mussel, *Mytilus edulis*, Meats Obtained From Cultivated and Natural Stock," Bohdan M. Slabyj, Donn L. Creamer, and Ruth H. True, 8:18-23.
 1322. "Catch-Effort and Price-Cost Trends in the Gulf of Mexico Shrimp Fishery: Implications on Mexico's Extended Jurisdiction," Vito Blomo, Wade L. Griffin, and John P. Nichols, 8:24-28.
 1323. "Estimating the Structure of Capacity Utilization in the Fishing Industry," Kenneth Ballard and Vito Blomo, 8:29-34.

SEPTEMBER

1324. "Composition of the Edible Portion of Raw (Fresh or Frozen) Crustaceans, Finfish, and Mollusks. III. Microelements," Virginia D. Sidwell, Audrey L. Loomis, Karen J. Loomis, Pauline R. Foncannon, and David H. Buzzell, 9:1-20.
 1325. "A Trawl Design: Employing Electricity to Selectively Capture Shrimp," Wilber R. Seidel and John W. Watson, Jr., 9:21-23.
 1326. "Use of Carbon Dioxide Dissolved in Refrigerated Brine for the Preservation of Pink Shrimp (*Pandalus* spp.)," Harold J. Barnett, Richard W. Nelson, Patrick J. Hunter, and Herman Groninger, 9:24-28.
 1327. "Status and Problems of the American Shrimp Industry," Roger W. Hutchinson, 9:29-31.

OCTOBER

1328. "Bacterial Diseases of Fishes and Their Control," S. F. Snieszko, 10:2-3.
 1329. "Diseases and Defense Mechanisms of the Lobster, *Homarus americanus*," James E. Stewart, 10:4.
 1330. "A New Bacterium (Presumptive *Vibrio* Species) Causing Ulcers in Flatfish," R. A. Robohm and C. Brown, 10:5-7.
 1331. "Bacteriologic Studies of Long Island Shellfish Hatcheries," Louis Leibovitz, 10:8.

1332. "Vibriosis in Maine and New Hampshire Salmonids," Evelyn S. Sawyer, 10:9.
 1333. "Anaerobic Bacteria as Possible Disease Agents in Fish," Lanny R. Udey, 10:10-11.
 1334. "Recent Development in Channel Catfish Virus Research," J. A. Plumb and P. R. Bowser, 10:12.
 1335. "Viral Diseases of the Blue Crab, *Callinectes sapidus*," Phyllis T. Johnson, 10:13-15.
 1336. "Piscine Erythrocytic Necrosis (PEN): A Viral Infection of the Atlantic Cod and Other Marine Fishes," Paul Reno, Marie Philippon-Fried, Bruce L. Nicholson, and Stuart Sherburne, 10:16-17.
 1337. "Viruses and Virulike Lesions in Marine Mollusks," C. Austin Farley, 10:18-20.
 1338. "Viruses and Virus Diseases of Salmonid Fishes," Philip E. McAllister, 10:21-23.
 1339. "Phosphonoacetic Acid Inhibition of Channel Catfish Virus Replication," Roger W. Koment, 10:24-25.
 1340. "Comments on Trends in Research on Parasitic Diseases of Shellfish and Fish," Victor Sprague, 10:26-29.
 1341. "Susceptibility Studies of Various Salmonids to Whirling Disease: Histological Staining and Spore Concentration Procedures," Joseph O'Grödnick, 10:30-31.
 1342. "Some Protozoan Diseases of Decapod Crustaceans," Thomas K. Sawyer and Sharon A. MacLean, 10:32-34.
 1343. "The Principal Parasitic Diseases of Warmwater Fishes," W. A. Rogers, 10:35-36.
 1344. "Trypanorhynch Infections in the Flesh of Sciaenid Fishes," Robin M. Overstreet, 10:37-38.
 1345. "Larval Nematodes Parasitic in Shellfish," Thomas C. Cheng, 10:39-42.
 1346. "Environmentally Related Diseases of Marine Fish and Shellfish," Carl J. Sindermann, 10:43.
 1347. "Nutrient Requirement of the Lobster and Nutrition Pathology," Robert C. Bayer, Margie Lee Gallagher, and Dale F. Leavitt, 10:44.
 1348. "Nutritionally Induced Cataracts in Salmonids Fed Purified and Practical Diets," Hugh A. Poston, Ronald C. Riis, Gary L. Rumsey, and H. George Ketola, 10:45.
 1349. "Environmental Parameters Affecting Fish Physiology in Water Reuse Systems," Thomas L. Meade, 10:46.
 1350. "PCB-Induced Alterations in Teleost Liver: A Model for Environmental Disease in Fish," David E. Hinton, James E. Klaunig, and Michael M. Lipsky, 10:47-50.
 1351. "A Review of Histopathological Effects of Selected Contaminants on Some Marine Organisms," George R. Gardner, 10:51-52.

NOVEMBER

1352. "Scallops and Their Utilization," John A. Peters, 11:1-9.
 1353. "Blue Crab Larval Culture: Methods and Management," Mark R. Millikin, 11:10-17.
 1354. "The 1976 Catch of Bowhead Whales, *Balaena mysticetus*, by Alaskan Eskimos," Willman M. Marquette, 11:18-27.

DECEMBER

1355. "Composition of the Edible Portion of Raw (Fresh or Frozen) Crustaceans, Finfish, and Mollusks. IV. Vitamins," Virginia D. Sidwell, Audrey L. Loomis, Pauline R. Foncannon, and David H. Buzzell, 12:1-16.
 1356. "Review of Oxygen Depletion and Associated Mass Mortalities of Shellfish in the Middle Atlantic Bight in 1976," Frank W. Steimle and Carl J. Sindermann, 12:17-26.

Editorial Guidelines for *Marine Fisheries Review*

Marine Fisheries Review publishes review articles, original research reports, significant progress reports, technical notes, and news articles on fisheries science, engineering, and economics, commercial and recreational fisheries, marine mammal studies, aquaculture, and U.S. and foreign fisheries developments. Emphasis, however, is on in-depth review articles and practical or applied aspects of marine fisheries rather than pure research.

Preferred paper length ranges from 4 to 12 printed pages (about 10-40 manuscript pages), although shorter and longer papers are sometimes accepted. Papers are normally printed within 4-6 months of acceptance. Publication is hastened when manuscripts conform to the following recommended guidelines.

The Manuscript

Submission of a manuscript to *Marine Fisheries Review* implies that the manuscript is the author's own work, has not been submitted for publication elsewhere, and is ready for publication as submitted. Commerce Department personnel should submit papers under completed NOAA Form 25-700.

Manuscripts must be typed (double-spaced) on high-quality white bond paper and submitted with two duplicate (but not carbon) copies. The complete manuscript normally includes a title page, a short abstract (if needed), text, literature citations, tables, figure legends, footnotes, and the figures. The title page should carry the title and the name, department, institution or other affiliation, and complete address (plus current address if different) of the author(s). Manuscript pages should be numbered and have 1½-inch margins on all sides. Running heads are not used. An "Acknowledgments" section, if needed, may be placed at the end of the text. Use of appendices is discouraged.

Abstract and Headings

Keep titles, heading, subheadings, and the abstract short and clear. Abstracts should be short (one-half page or less) and

double-spaced. Paper titles should be no longer than 60 characters; a four- to five-word (40 to 45 characters) title is ideal. Use heads sparingly, if at all. Heads should contain only 2-5 words; do not stack heads of different sizes.

Style

In style, *Marine Fisheries Review* follows the "U.S. Government Printing Office Style Manual." Fish names follow the American Fisheries Society's Special Publication No. 6, "A List of Common and Scientific Names of Fishes from the United States and Canada," third edition, 1970. The "Merriam-Webster Third New International Dictionary" is used as the authority for correct spelling and word division. Only journal titles and scientific names (genera and species) should be italicized (under-scored). Dates should be written as 3 November 1976. In text, literature is cited as Lynn and Reid (1968) or as (Lynn and Reid, 1968). Common abbreviations and symbols such as mm, m, g, ml, mg, and °C (without periods) may be used with numerals. Measurements are preferred in metric units; other equivalent units (i.e., fathoms, °F) may also be listed in parentheses.

Tables and Footnotes

Tables and footnotes should be typed separately and double-spaced. Tables should be numbered and referenced in text. Table headings and format should be consistent; do not use vertical rules.

Literature Citations

Title the list of references "Literature Citations" and include only published works or those actually in press. Citations must contain the complete title of the work, inclusive pagination, full journal title, the year and month and volume and issue numbers of the publication. Unpublished reports or manuscripts and personal communications must be footnoted. Include the title, author, pagination of the manuscript or report, and the address where it is on file. For personal communications, list the name, affiliation, and address of the communicator.

Citations should be double-spaced and listed alphabetically by the senior author's surname and initials. Co-authors should be listed by initials and surname. Where two or more citations have the same author(s), list them chronologically; where both author and year match on two or more, use lower-case alphabet to distinguish them (1969a, 1969b, 1969c, etc.).

Authors must double-check all literature cited; they alone are responsible for its accuracy.

Figures

All figures should be clearly identified with the author's name and figure number, if used. Figure legends should be brief and a copy may be taped to the back of the figure. Figures may or may not be numbered. Do not write on the back of photographs. Photographs should be black and white, 8-× 10- inches, sharply focused glossies of strong contrast. Potential cover photos are welcome but their return cannot be guaranteed. Magnification listed for photomicrographs must match the figure submitted (a scale bar may be preferred).

Line art should be drawn with black India ink on white paper. Design, symbols, and lettering should be neat, legible, and simple. Avoid freehand lettering and heavy lettering and shading that could fill in when the figure is reduced. Consider column and page sizes when designing figures.

Finally

First-rate, professional papers are neat, accurate, and complete. Authors should proofread the manuscript for typographical errors and double-check its contents and appearance before submission. Mail the manuscript flat, first-class mail, to: Editor, *Marine Fisheries Review*, Scientific Publications Office, National Marine Fisheries Service, NOAA, 1107 N.E. 45th Street, Room 450, Seattle, WA 98105.

The senior author will receive 50 reprints (no cover) of his paper free of charge and 100 free copies are supplied to his organization. Cost estimates for additional reprints can be supplied upon request.

UNITED STATES
DEPARTMENT OF COMMERCE

NATIONAL OCEANIC AND ATMOSPHERIC ADMINISTRATION
NATIONAL MARINE FISHERIES SERVICE
SCIENTIFIC PUBLICATION OFFICE
MARINE FISHERIES REVIEW
ROOM 450

1107 N.E. 45TH ST.
SEATTLE, WA 98105
OFFICIAL BUSINESS

POSTAGE AND FEES PAID
U.S. DEPARTMENT OF COMMERCE
COM-210

Controlled Circulation Rate



MFR UNIVM300UF155DUE004R 1
UNIV MICROFILMS INTL
SERIALS PROCESSING
300 N ZEEB RD
ANN ARBOR MI 48106

